

Diffusive behavior for randomly kicked Newtonian particles in a spatially periodic medium

Jeremy Clark and Christian Maes

Instituut voor Theoretische Fysica
K.U.Leuven, Belgium

September 14, 2010

Abstract

We prove a central limit theorem for the momentum distribution of a particle undergoing an unbiased spatially periodic random forcing at exponentially distributed times without friction. The start is a linear Boltzmann equation for the phase space density, where the average energy of the particle grows linearly in time. Rescaling time, the momentum converges to a Brownian motion, and the position is its time-integral showing superdiffusive scaling with time $t^{3/2}$. The analysis has two parts: (1) to show that the particle spends most of its time at high energy, where the spatial environment is practically invisible; (2) to treat the low energy incursions where the motion is dominated by the deterministic force, with potential drift but where symmetry arguments cancel the ballistic behavior.

1 Introduction

Recent times show a renewed great interest in obtaining diffusive behavior from microscopically defined dynamics. The motivation is much older, to derive, as Fourier, Navier or Boltzmann first did in their ways and times, irreversible and dissipative behavior starting from the reversible microscopic laws. The limiting behavior is often associated to a conserved quantity like energy in classical mechanics and the challenge is then to express the (energy) current in terms of gradients of the (energy) density itself. Obviously, for the sharpness of the limit, some scaling must be done, combined with typicality arguments on the level of the initial or boundary conditions. For example, more recently the search for a rigorous derivation of Fourier's law of heat conduction was relaunched in [2] and many attempts and models have been taken up after that.

More modestly one starts half-way with an effective description on the level of single-particle dynamics. The one particle phase space density then really refers to a cloud of weakly interacting particles brought in contact with some environment, and the conserved quantity is simply the number of particles. In the present paper we study the diffusive scaling limit of a massive particle in a one-dimensional periodic potential to which we add random forcing. The latter is not derived from first principles, but has various physical motivations. Heuristically, the forcing corresponds to the random collisions with an effectively infinite temperature granular bath. The granular structure is in the discrete kicks the particle undergoes at random exponentially

distributed times. Depending on its present position, the distribution of the momentum kicks differs. Together with the potential that specifies the spatial inhomogeneity and makes the problem nontrivial.

The spatial heterogeneity brings us to a second motivation of the present work: the study of active particles where flights of ballistic motion are interrupted by spatially depending re-orientations, or the statistical characterization of particle trajectories in active heterogeneous fluids, see e.g. [15]. Our present work adds a rigorous result establishing under what conditions diffusive behavior for the momentum and superdiffusive behavior for the position get realized.

Finally, the present contribution fits in the long tradition of proofs of the central limit theorem (invariance principle, and its modifications) for additive functionals, position as the integral of the momenta and momenta as the integral of the forces. Much of all that for studies of interacting particle systems started from the pioneering work in [12]. In the spirit of the present work the papers [4, 5] added very important symmetry considerations, making it possible to apply the work to strongly dependent variables. The fact that these arguments avoided the use of mixing assumptions or strong enough decay of time-correlations appears like an important lesson for today's pursuit of diffusive behavior mentioned at the very beginning of this introduction. Indeed, one often emphasizes that strong enough chaoticity assumptions are needed in the mathematical control of the transition from the microscopic reversible laws to macroscopic irreversible behavior. One then refers for example to the problem of obtaining regular transport properties via well-controlled Green-Kubo expressions where some temporal decay certainly seems necessary. These Green-Kubo relations are however needed only for very special observables, and not for all possible even microscopically defined quantities. It is therefore very welcome if symmetry considerations can help to establish diffusive behavior for certain classes of functionals that share symmetry properties with the relevant observables of statistical mechanics. The present paper is not starting from the microscopic classical mechanical world, but it does deal with the problem of exploiting symmetry to cancel ballistic behavior. Other more recent work on the central limit theorem that shares important ambitions with the present study includes [10, 9, 1, 6]. For the study of fluctuations in Markov processes with an overview of central limit results, we refer to the recent book [13].

The next section introduces the model, the results and the main strategy of the proof. The momentum variable is not autonomous since it is coupled to the position of the particle. Its changes come from two sources, the momentum jumps by the external Poisson noise and the acceleration due to the presence of the potential. That is translated in the structure of the argument. The idea is to obtain a martingale central limit theorem for the momentum jumps while the effect of the potential should vanish in the long time limit. Section 3 establishes that, most of the time, the particle's energy grows linearly with time. That is sufficient to show in Section 4 that the absolute value of the momentum process converges in distribution to the absolute value of a Brownian motion. Next, in Section 5, follow the estimates characterizing the motion at high energy, where the (bounded) potential has very little effect. The low energy motion is discussed under section 6. There the drift due to the potential gets controlled by symmetry arguments. The combination of high and low energy estimates yields the main result of Section 7.

2 Main result

2.1 Informal description

Consider a one-dimensional classical particle whose position and momentum (X_t, K_t) evolve deterministically with Hamiltonian $H(x, k) = \frac{1}{2}k^2 + V(x)$ for some bounded periodic potential $0 \leq V(x) \leq \bar{V}$ except at Poisson times at which the particle may receive a momentum kick from the environment. That is, independent of its current momentum k and at the rate $j_x(v)$ when its current position is x the particle receives a momentum jump v . On the level of the phase space densities, the dynamics we consider is then governed by the linear Boltzmann equation

$$\frac{d}{dt}P_t(x, k) = -k \frac{\partial P_t}{\partial x}(x, k) + \frac{dV}{dx}(x) \frac{\partial P_t}{\partial k}(x, k) + \int_{\mathbb{R}} dv j_x(v) (P_t(x, k-v) - P_t(x, k)) \quad (2.1)$$

for the phase space probability density $P_t(x, k) \in L^1(\mathbb{R}^2)$ for the particle at each time $t \geq 0$. The rates j_x have the same periodicity as the potential V . It is however probabilistically simpler to imagine a universal Poisson clock having rate $\mathcal{R} > 0$, such that when the alarm rings, a (biased) coin is tossed to decide whether or not a momentum kick will occur. The probability of the coin and the distribution of the momentum jump v are respectively $0 \leq \kappa(a) \leq 1$ and $\mathcal{P}_a(v)$, where $a = X_t \bmod 1$ is the position (modulo the period 1 of the potential V) at the Poisson time t . We assume that the momentum jumps are symmetric $\mathcal{P}_a(-v) = \mathcal{P}_a(v)$ and that there is a uniform lower bound for the coin probabilities $0 < \nu \leq \kappa(a)$. Then, in (2.1), $j_x(v) = \mathcal{R} \kappa(x) \mathcal{P}_x(v)$.

Our main result is to show, under certain technical conditions, that the normalized variables $(t^{-\frac{3}{2}}X_{st}, t^{-\frac{1}{2}}K_{st})$, $s \geq 0$ approach the process $(\int_0^s dr \mathbf{B}_r, \mathbf{B}_s)$ in distribution where $\mathbf{B}_s$ is Brownian motion whose diffusion constant σ depends on the spatial average of the periodic noise $\sigma = \int_0^1 da \int_{\mathbb{R}} dv j_a(v) v^2$.

There is clearly no energy relaxation in (2.1), since, no matter where you start, the time-derivative of the expected energy satisfies

$$\frac{d}{dt}\mathbb{E}[E_t] = \frac{d}{dt} \int_{\mathbb{R}^2} dx dk \left(\frac{k^2}{2} + V(x) \right) P_t(x, k) = \frac{1}{2} \int_{\mathbb{R}^2} dx dk \left(\int_{\mathbb{R}} dv j_x(v) v^2 \right) P_t(x, k) \quad (2.2)$$

and thus the mean energy grows linearly as

$$\mathbb{E}[E_0] + \frac{t}{2} \inf_{a \in [0, 1]} \int_{\mathbb{R}} dv j_a(v) v^2 \leq \mathbb{E}[E_t] \leq \mathbb{E}[E_0] + \frac{t}{2} \sup_{a \in [0, 1]} \int_{\mathbb{R}} dv j_a(v) v^2.$$

Moreover, as we will show, by time t not only the average but also the typical energy is of order t . Since the potential $V(x)$ is bounded, the absolute value of the momentum is then $|k| \propto t^{\frac{1}{2}}$. As a consequence of having high momentum, the particle will pass through one period of the potential (periodic cell) much faster than the time scale of the Poisson clock governing the noise. The particle then effectively “feels” a spatial average of the noise in which the averaged distribution of a jump $\tilde{P}(v)$ and the averaged Poisson rate are $\tilde{\mathcal{R}}$

$$\tilde{P}(v) = \frac{\int_0^1 da \kappa(a) \mathcal{P}_a(v)}{\int_0^1 da \kappa(a)} \quad \text{and} \quad \tilde{\mathcal{R}} = \mathcal{R} \int_0^1 da \kappa(a)$$

Moreover, at very high momentum, the force field by the potential can only displace the momentum by relatively negligible values. The effective dynamics at high energy is thus

$$\frac{d}{dt}P_t(x, k) = -k \frac{\partial P_t}{\partial x}(x, k) + \tilde{\mathcal{R}} \int_{\mathbb{R}} dv \tilde{P}(v) (P_t(x, k - v) - P_t(x, k)) \quad (2.3)$$

This dynamics is translation invariant and so the momentum process has now become a Markov process. Showing that $(t^{-\frac{3}{2}}X_{st}, t^{-\frac{1}{2}}K_{st})$ converges to $(\int_0^s dr \mathbf{B}_r, \mathbf{B}_s)$ is then straightforward.

Another way of expressing this result on the level of single-time marginals is to consider the rescaled density $t^2 P_t(t^{3/2}x, t^{1/2}k)$ at time t . Its limit $t \uparrow \infty$ is Gaussian $P_\infty(x, k)$,

$$P_\infty(x, k) = \frac{\sqrt{3}}{\pi \sigma} e^{-\frac{6}{\sigma}(x - \frac{k}{2})^2 - \frac{1}{2\sigma}k^2}$$

and $\sigma = \int_0^1 dx \int_{\mathbb{R}} dv j_x(v) v^2$. The coupling between position x and momentum k results from the correlation between the Brownian motion (for the momentum) and its time-integral (for the position).

2.2 Strategy of proof

The process (X_t, K_t) is Markovian over the set of right-continuous paths (having left limits) from $t \in \mathbb{R}^+$ to $\mathbb{R}^2$, bounded over finite time intervals. Since the position variable is an integral of the momentum, the proof that $(t^{-\frac{3}{2}}X_{st}, t^{-\frac{1}{2}}K_{st})$ converges to $(\int_0^s dr \mathbf{B}_r, \mathbf{B}_s)$ for a Brownian motion $\mathbf{B}_s$ is implied by showing that the momentum component converges to a Brownian motion. The momentum process can be written as

$$K_{st} = K_0 + M_{st} + \int_0^{st} dr \frac{dV}{dx}(X_r) \quad (2.4)$$

where M_t is the martingale of jumps, $M_t = \sum_s^t v_s$ over the jump times $0 \leq s \leq t$ in the Poisson process at rate $\mathcal{R}$ and v_s is the actual momentum kick. On the other hand, $\int_0^t dr \frac{dV}{dx}(X_r)$ is the net drift due to the conservative force up to time t .

The analysis splits into two semi-independent parts corresponding to the two last terms in (2.4). First we show that the momentum jump part $t^{-\frac{1}{2}}M_{st}$ converges to a Brownian motion. That requires establishing a martingale central limit theorem. Because of the inhomogeneity in the momentum jumps we need to prove that there is asymptotic regularity in the variances of the momentum jumps (quadratic variation process). That is realized because the particle spends most of its time at high energy where translation invariance is recovered. We call that the high energy analysis.

Secondly, for the low energy analysis we show that the drift process $t^{-\frac{1}{2}} \int_0^{st} dr \frac{dV}{dx}(X_r)$ makes a vanishing contribution for large times; in other words the variance of the time integral of the drift converges to zero. Indeed note again that the final process, obtained after scaling, does not depend on the potential V . The superdiffusion of the position makes the position process almost deterministic and the random kicks become rare compared to the fast movement of the particle. Mathematically, in section 6 we use that periods of low energy are well separated by times of high energy. There is therefore some independence between the low energy incursions. Moreover symmetry arguments constrain the gained momentum in each such incursion to have zero expectation.

2.3 Main theorem

Our main mathematical result is a central limit theorem for the momentum process. We give here the precise statement.

Assumptions List 2.1.

- I. There exists $0 < r_1$ such that for all $a \in [0, 1]$, $r_1 \leq \int_{\mathbb{R}} dv j_a(v) v^2$.*
- II. There is $\rho > 0$ such that for all $a \in [0, 1]$, $\int_{\mathbb{R}} dv \mathcal{P}_a(v) v^4 \leq \rho$.*
- III. $j_a(v) = j_a(-v)$*
- IV. $\bar{V} > V(x) \geq 0$ is bounded and has a bounded derivative.*

The first three assumptions are on the rate of momentum jumps. They should be symmetric, allow spreading but still have a fourth momentum. For the Hamiltonian part, both the potential and the force is bounded. The assumptions of List 2.1 are designed to be the minimal assumptions for Section 3, and most results from later sections require both List 2.1 and some of the assumptions from List 2.2.

Assumptions List 2.2.

- i. There exists $\mathcal{C}$ and $\eta > 0$ such that for all $a \in [0, 1]$ and $v, w \in \mathbb{R}$ with $|v| - |w| \geq 0$*

$$\mathcal{P}_a(w) \leq \mathcal{C} e^{-\eta(|v|-|w|)} \mathcal{P}_a(v).$$

- ii. There exists a μ such that for all $a \in [0, 1]$,*

$$\sup_{v \in \mathbb{R}} \left[(\mathcal{P}_a(v))^{-1} (1 + |v|)^{-1} \sup_{|w-v| \leq 1} \left| \frac{d\mathcal{P}_a}{dv}(w) \right| \right] \leq \mu.$$

- iii. There exists a reflection R on the torus such that $V(R(x)) = V(x)$ and $j_{R(a)}(v) = j_a(v)$ for $a \in [0, 1]$ and $v \in \mathbb{R}$.*

Condition (i) implies that the Laplace transform of $\mathcal{P}_a$ is finite in a neighborhood around zero and thus that the fourth moment as in (II) and all other moments are finite.

In later sections, r_1 , r_2 , ν will be defined as

$$r_1 = \inf_{a \in [0,1]} \int_{\mathbb{R}} dv j_a(v) v^2, \quad r_2 = \sup_{a \in [0,1]} \int_{\mathbb{R}} dv j_a(v) v^2, \quad \nu = \inf_{a \in [0,1]} \kappa_a.$$

Since $j_a(v) = \mathcal{R} \kappa_a \mathcal{P}_a(v)$, the condition (I) and (II) with Jensen's inequality imply that $0 < r_1 \mathcal{R}^{-1} \rho^{-\frac{1}{2}} \leq \nu$. Also by (II) and Jensen's inequality, $r_2 < \mathcal{R} \rho^{\frac{1}{2}} < \infty$.

(III) and (iii) are the symmetries that we assume for the dynamics. (III) says that for every point $a \in [0, 1]$ in the periodic cell, the rate of kicks by a momentum v occurs with the same rate as kicks by a momentum $-v$. (iv) specifies that in addition to the periodicity of the dynamics, there is also a spatial reflection symmetry. The combination of these symmetries forms a “momentum time-reversal symmetry” which is used in Section 6.2.

(ii) is a technical assumption so that the values of the derivative $|\frac{d\mathcal{P}_a}{dv}(w)|$ for w in a neighborhood around v cannot be too large compared to the value $\mathcal{P}_a(v)$. The constraint becomes more flexible at large $|v|$, where the ratio is allowed to increase as $|v|$. This condition effectively forbids densities with tails that vanish faster than a Gaussian density $G(v)$ in which

$$\frac{|\frac{dG}{dv}(v)|}{G(v)} \propto |v|.$$

The support of $\mathcal{P}_a(v)$ for each a cannot be finite for instance. Avoiding this decay of $\mathcal{P}(v)$ as $|v| \rightarrow \infty$ is not essential to the analysis, but generalizing the condition (ii) (for instance by replacing $(1 + |v|)$ by $(1 + |v|)^m$) requires making other conditions more complicated.

Theorem 2.3 (Main result). *Assume List 2.1, List 2.2, and that the initial joint phase space distribution $P_0(x, k) \in L^1(\mathbb{R}^2)$ has finite second moments. In the limit $t \rightarrow \infty$, $(t^{-\frac{3}{2}}X_{st}, t^{-\frac{1}{2}}K_{st})$ converges in distribution to $(\int_0^s dr \mathbf{B}_r, \mathbf{B}_s)$ where $\mathbf{B}_s$ is Brownian motion with diffusion constant $\sigma = \int_0^1 da \int_{\mathbb{R}} dv j_a(v) v^2$.*

3 A martingale central limit theorem

In this section, we prove that the typical energy for the particle is on the order of t . That implies a regularity in the momentum process, at least concerning its absolute value and for the quadratic variation of the momentum jumps. The net result is a martingale central limit theorem for the martingale part in (2.4). Note that we always assume the natural filtration $\mathcal{F}_t$ specifying the Markov process up to time t .

Theorem 3.1. *Assume List 2.1 and (i)-(ii) of List 2.2 and that the initial joint distribution $P_0(x, k)$ has finite second moments. Then $t^{-\frac{1}{2}}M_{st}$ converges in distribution to a Brownian motion $\mathbf{B}_s$ with diffusion constant σ .*

The proof follows in Section 7. It will be built on the lemma's below and in the next section.

The following lemma relies on a martingale central limit theorem [7] and on having some bounds for the time-change of that central limit theorem. The result applies to more general class of martingales, but we develop it here to the martingale process

$$M'_t = \sum_{n=1}^{\mathcal{N}_t} w_n S(K_{t_n^-}),$$

where t_n are the Poisson time for the underlying Poisson clock $\mathcal{N}_t$ with rate $\mathcal{R}$, $S(K_{t^-})$ is the left limit up to time t for the sign of the momentum, and $w_n = M_{t_n} - M_{t_n^-}$. Note that w_n is zero if at the Poisson time t_n there is no momentum jump, and it is equal to the momentum jump if it does happen.

Lemma 3.2. *For $T_\epsilon^{(t)} = \int_0^1 dr \chi(|t^{-\frac{1}{2}}M'_{rt}| > \epsilon)$, there exists a $C > 0$ such that for all $\epsilon, \delta > 0$*

$$\liminf_{t \rightarrow \infty} \Pr[T_\epsilon^{(t)} \geq 1 - \delta] \geq 1 - C \frac{r_2^{\frac{1}{2}} \epsilon}{r_1 \delta}.$$

The same result remains true with $|t^{-\frac{1}{2}}M'_{st}|$ replaced by $t^{-\frac{1}{2}}M'_{st} - \inf_{0 \leq u \leq s} t^{-\frac{1}{2}}M'_{ut}$.

Proof. We start from the lower bound:

$$\begin{aligned} \Pr[T_\epsilon^{(t)} \leq 1 - \delta] &= 1 - \Pr\left[\int_0^1 dr \chi(t^{-\frac{1}{2}}|M'_{st}| \leq \epsilon) \geq \delta\right] \\ &\geq 1 - \frac{1}{\delta} \mathbb{E}\left[\int_0^1 ds \chi(|t^{-\frac{1}{2}}M'_{st}| \leq \epsilon)\right] = 1 - \frac{1}{\delta} \int_0^1 ds \Pr[|t^{-\frac{1}{2}}M'_{st}| \leq \epsilon] \end{aligned} \quad (3.1)$$

Define $\tilde{B}_u^{(t)} = t^{-\frac{1}{2}}M'_{\tau_u t}$, where τ_u is the hitting time

$$\tau_u = \inf\left\{s \geq 0 \mid \frac{1}{r_1 t} \langle M' \rangle_{st} \geq u\right\},$$

and $\langle M' \rangle_t$ is the predictable quadratic variation of M' up to time t . In our situation, $\langle M' \rangle_t$ has the form

$$\langle M' \rangle_t = \int_0^t dr \int_{\mathbb{R}} dv j_{X_r}(v) v^2.$$

By the martingale central limit theorem, $\tilde{B}_u^{(t)}$ converges to a Brownian motion in the uniform metric. The Lindberg condition is guaranteed by the boundedness of the fourth moment of single momentum jumps in (II) of List 2.1 and the fact that the jump times occur according to a Poisson clock (having rate $\mathcal{R}$).

We also have $t^{-\frac{1}{2}}M'_{st} = \tilde{B}_{R_s}^{(t)}$, where $R_s = \frac{1}{r_1 t} \langle M' \rangle_{st}$, since τ_u and R_s are inverses of one another. By (II) of List 2.1 and the discussion following it, $\int_{\mathbb{R}} dv j_a(v) v^2$ ranges between the values $0 < r_1 \leq r_2$ for $a \in [0, 1]$. It follows that,

$$u \frac{r_1}{r_2} \leq \tau_u \leq u \quad \text{and} \quad s \leq R_s \leq s \frac{r_2}{r_1}. \quad (3.2)$$

By (3.2), R_s has the range $s \leq R_s \leq \frac{r_2}{r_1} s$, and thus

$$\int_0^1 ds \chi(|t^{-\frac{1}{2}}M'_{st}| \leq \epsilon) = \int_0^1 ds \chi(|\tilde{B}_{R_s}^{(t)}| \leq \epsilon) \leq \int_0^{\frac{r_2}{r_1}} du \chi(|\tilde{B}_u^{(t)}| \leq \epsilon).$$

Taking the expectation of the right-hand side and using the fact that $\tilde{B}_s^{(t)}$ approaches a Brownian motion, we have that

$$\mathbb{E}\left[\int_0^{\frac{r_2}{r_1}} du \chi(|\tilde{B}_u^{(t)}| \leq \epsilon)\right] = \int_0^{\frac{r_2}{r_1}} du \Pr[|\tilde{B}_u^{(t)}| \leq \epsilon] \longrightarrow \int_0^{\frac{r_2}{r_1}} du \int_{-\epsilon}^{\epsilon} dx \frac{e^{-\frac{x^2}{2r_1 s}}}{\sqrt{2\pi r_1 s}}$$

By a change of variables, the right side is bounded by a constant multiple of $\epsilon r_1^{-1} r_2^{\frac{1}{2}}$. With (3.1) this proves the result.

To generalize the result to $H_s^{(t)} = t^{-\frac{1}{2}}M'_{st} - \inf_{0 \leq a \leq s} t^{-\frac{1}{2}}M'_{at}$, we make the same time-change τ , to define a process $Z_u^{(t)} = H_{\tau_u}^{(t)}$. Since $\tilde{B}_u^{(t)} = t^{-\frac{1}{2}}M'_{\tau_u t}$ converges to a Brownian motion, it will follow that $Z_u^{(t)}$ converges to the absolute value of a Brownian motion. Indeed, the function $f : L^\infty([0, 1]) \rightarrow L^\infty([0, 1])$ defined by $f(x_s) = x_s + \sup_{0 \leq r \leq s} -x_r$ (read the supremum as an essential supremum) satisfies $\|f(x_s) - f(y_s)\|_\infty \leq 2\|x_s - y_s\|_\infty$. Thus the convergence of $\tilde{B}_u^{(t)}$ to a Brownian motion $\mathbf{B}_u$ implies that $f(\tilde{B}_u^{(t)})$ converges in distribution to $f(\mathbf{B}_u)$. However, by the basic result for Brownina motion [11], $f(\mathbf{B}_u)$ is equal in distribution to $|\mathbf{B}_u|$. Thus we can apply the same reasoning as above to get the result. $\square$

The lemma below will be used in the proof of Lemma 3.4. Its proof follows from basic calculus but requires consideration of several cases.

Lemma 3.3. *For $\bar{V} = \sup_{x \in \mathbb{R}} V(x) < \infty$, there exists a $c > 0$ such that*

$$\begin{aligned} & \left| 2^{\frac{1}{2}} w S(k) - \left[\left(\frac{1}{2} |k+w|^2 + V \right)^{\frac{1}{2}} - \left(\frac{1}{2} |k-w|^2 + V \right)^{\frac{1}{2}} \right] \right|^2 \\ & \leq 2 w^2 1_{|w| > J} + c J \left[\left(\frac{1}{2} |k+w|^2 + V \right)^{\frac{1}{2}} + \left(\frac{1}{2} |k-w|^2 + V \right)^{\frac{1}{2}} - 2 \left(\frac{1}{2} k^2 + V \right)^{\frac{1}{2}} \right] \end{aligned} \quad (3.3)$$

for all $k, w \in \mathbb{R}$, $0 \leq V \leq \bar{V}$, and $J \geq V^{\frac{1}{2}}$.

Proof. When $V = 0$, we use the identity

$$\left| S(k)w - \frac{1}{2}(|k+w| - |k-w|) \right| = \frac{1}{2}(|k+w| + |k-w| - 2|k|).$$

Hence, for $c = 2$,

$$\left| 2^{\frac{1}{2}} S(k)w - 2^{-\frac{1}{2}}(|k+w| - |k-w|) \right|^2 \leq 2 v^2 \chi(|w| > J) + \frac{c J}{2}(|k+w| + |k-w| - 2|k|), \quad (3.4)$$

since $|S(k)w - \frac{1}{2}(|k+w| - |k-w|)| \leq |w|$.

When $V \neq 0$, we define

$$\begin{aligned} M(w, k) &= 2 w S(k) - (|k+w|^2 + 1)^{\frac{1}{2}} + (|k-w|^2 + 1)^{\frac{1}{2}} \\ N(w, k) &= (|k+w|^2 + 1)^{\frac{1}{2}} + (|k-w|^2 + 1)^{\frac{1}{2}} - 2(k^2 + 1)^{\frac{1}{2}} \end{aligned}$$

We can divide the inequality (3.3) by V , and let $2^{-\frac{1}{2}} V^{-\frac{1}{2}} k \rightarrow k$, $2^{-\frac{1}{2}} V^{-\frac{1}{2}} w \rightarrow v$, and $J V^{-\frac{1}{2}} \rightarrow J$ so that the inequality we must show is

$$|M(w, k)|^2 \leq 4 w^2 1_{|w| > J} + c J N(w, k) \quad (3.5)$$

for some c . For $|w| > J$ the inequality is trivial, since the $M(w, k)$ is smaller than $2|w|$. Hence we can take $J = |w| \vee 1$. If we show that there is such a c' such that (3.5) holds for $(w, k) \in \mathbb{R}^2 - D$ for the compact domain $D = ([-L, -L^{-1}] \cup [L^{-1}, L]) \times [-L, L]$ for some $L > 1$, then, since $M(w, k)$ is bounded inside D and $N(w, k)$ is bounded away from zero inside of D , it follows that there exists a c verifying (3.5) for all $(w, k) \in \mathbb{R}^2$. The proof that there exists such a c' for all $(w, k) \in \mathbb{R}^2 - D$ for some $L \gg 1$ is partitioned into the following main cases

1. $|w| \ll 1$,
2. $|w| \gg 1$ and $|k| \geq |w| + |w|^{\frac{1}{2}}$,
3. $|w| \gg 1$ and $|k+w| \leq |w|^{\frac{1}{2}}$ or $|k-w| \leq |w|^{\frac{1}{2}}$,
4. $|w| \gg 1$ and $|k| \leq |w|^{\frac{1}{2}}$.

We go through each case and we show that $|M(w, k)|^2$ is smaller than some constant multiple of $(|w| \vee 1) N(w, k)$.

Case (1): The origin can only be between $k - w$ and $k + w$ when $|k| \leq |w| \ll 1$, in which case

$$|M(w, k)|^2 \leq w^2 < 2w^2 = 4\left[\frac{1}{2}(k + w)^2 + \frac{1}{2}(k - w)^2 - k^2\right] \approx 4N(w, k).$$

When the origin is not between $k - w$ and $k + w$ then by Taylor's formula for $(|k + w|^2 + 1)^{\frac{1}{2}}$ up to first order and using that $k + a$ has the same sign as k for $|a| \leq |w|$,

$$M(w, k) = S(k) \int_{-w}^w da \frac{|k + a| - (|k + a|^2 + 1)^{\frac{1}{2}}}{(|k + a|^2 + 1)^{\frac{1}{2}}}. \quad (3.6)$$

Using Taylor's formula to second order,

$$N(w, k) = \int_{-|w|}^{|w|} da \frac{|w| - |a|}{(|k + a|^2 + 1)^{\frac{3}{2}}}. \quad (3.7)$$

for all w and k . From these formulas we can take upper and lower bounds for $|M(w, k)|^2$ and $N(w, k)$ respectively:

$$|M(w, k)|^2 \leq |w|^2 \frac{||k| - |w| - ((|k| - |w|)^2 + 1)^{\frac{1}{2}}|^2}{(|k| - |w|)^2 + 1} \leq |w|^2 \frac{1 \wedge (|k| - |w|)^{-2}}{(|k| - |w|)^2 + 1}$$

$$N(w, k) \geq \frac{1}{2}|w|^2 \frac{1}{((|k| + |w|)^2 + 1)^{\frac{3}{2}}},$$

where we have used the monotonicity of the functions involved, and the second inequality for $|M(w, k)|^2$ includes the inequality

$$||a| - (a^2 + 1)^{\frac{1}{2}}| \leq 1 \wedge |a|^{-1}. \quad (3.8)$$

For large $|k|$, $|M(w, k)|^2$ has order $|k|^{-4}$ and $N(w, k)$ has order $|k|^{-3}$. Thus we can find a constant c' so that (3.5) holds for all (w, k) with $|w| < L^{-1}$ for some $L > 1$.

Case (2): When $|k| \geq |w| + |w|^{\frac{1}{2}}$, then we can apply (3.6), (3.7), and the same estimates as above to show that the asymptotics for $|M(w, k)|^2$ has lower order than $|v| N(v, k)$.

Case (3): It is convenient to follow the pattern in (3.4) and write $M(w, k)$ as $M = M_1 + M_2$, where

$$M_1(w, k) = 2v s(k) + |k + w| - |k - w|$$

$$M_2(w, k) = (|k + w|^2 + 1)^{\frac{1}{2}} - |k + w| - (|k - w|^2 + 1)^{\frac{1}{2}} + |k - w|.$$

Notice that $M_1(w, k)$ will only be non-zero when $|k| \leq |w|$, and that it can be written as $M_1(w, k) = 2S(k)S(w)(|w| - |k|)\chi(|w| \geq |k|)$. Applying the inequality $|x + y|^2 \leq 2x^2 + 2y^2$:

$$|M(w, k)|^2 \leq 2M_1^2(w, k) + 2M_2^2(w, k). \quad (3.9)$$

We start by bounding M_1 , since it will be used for the M_2 term. For $|w| \gg 1$ we claim that

$$(|w| - |k|)\chi(|w| \geq |k|) < N(w, k), \quad (3.10)$$

which implies a bound for M_1 , since

$$M_1 = 4(|w| - |k|)^2 \chi(|w| \geq |k|) \leq 4|w|(|w| - |k|) \chi(|w| \geq |k|) \leq 4|w| M_2(w, k).$$

The expression $(|w| - |k|) \chi(|w| \geq |k|)$ has its maximum at $k = 0$ and decreases linearly to zero at $|k| = |w|$. At $k = 0$, $(|w| - |k|) \chi(|w| \geq |k|) = |w| < 2|w| \approx N(0, k)$. However, $N(w, k)$ decreases at a slightly less than linear rate and $N(w, w) \approx 1$.

For the $M_2^2(w, k)$ term, we will finally use the case condition that either $|k + w| \leq |w|^{\frac{1}{2}}$ or $|k - w| \leq |w|^{\frac{1}{2}}$. Without loss of generality, let it be that $|k - w| \leq |w|^{\frac{1}{2}}$. Then $|k + w| \gg 1$ and so by (3.8) $\left| (|k + w|^2 + 1)^{\frac{1}{2}} - |k + w| \right| \sim |k + w|^{-1} \sim |w|^{-1}$. Thus

$$\begin{aligned} & \left((|k + w|^2 + 1)^{\frac{1}{2}} - |k + w| - (|k - w|^2 + 1)^{\frac{1}{2}} + |k - w| \right)^2 \\ &= \left[(|k - w|^2 + 1)^{\frac{1}{2}} - |k - w| \right]^2 + O(|w|^{-1}), \end{aligned} \quad (3.11)$$

which is $O(1)$. On the other hand,

$$|w| N(w, k) \approx |w| \left[(|k - w|^2 + 1)^{\frac{1}{2}} + |k + w| - |k| + O(|w|^{-1}) \right] = O(|w|^2),$$

since $(|k - w|^2 + 1)^{\frac{1}{2}} = O(|w|^{\frac{1}{2}})$. Thus $|w| N(w, k)$ has higher order than (3.11).

Case (4): Notice that

$$M_2(w, k) = \int_{-w}^w da \frac{k + a - S(k + a) (|k + a|^2 + 1)^{\frac{1}{2}}}{(|k + a|^2 + 1)^{\frac{1}{2}}}$$

where we have expanded the term $(|k + w|^2 + 1)^{\frac{1}{2}}$ appearing in M_2 to first order as in case (1) and we have written $|k + w| - |k - w| = \int_{-w}^w da S(k + a)$. Now,

$$\begin{aligned} & \left| \int_{-w}^w da \frac{k + a - S(k + a) (|k + a|^2 + 1)^{\frac{1}{2}}}{(|k + a|^2 + 1)^{\frac{1}{2}}} \right| \\ & \leq 2(|w| - |k|) + \left| \int_{-w}^w da \chi(|k + a| \geq |w|) \frac{k + a - S(k + a) (|k + a|^2 + 1)^{\frac{1}{2}}}{(|k + a|^2 + 1)^{\frac{1}{2}}} \right| \\ & \leq 2(|w| - |k|) + 2 \int_{|w|^{\frac{1}{2}}}^{|w|} da |a|^{-2} = 2(|w| - |k|) + O(|w|^{-\frac{1}{2}}), \end{aligned} \quad (3.12)$$

where in the first inequality we have used that the integrand on the left is bounded by 1 over the interval $a \in [k - 2|w|, k + 2|w|]$. The second inequality follows since $|k + a|^{-2}$ bounds the integrand, and $\int_{|w|^{\frac{1}{2}}}^{|w|} da |a|^{-2}$ is the smallest that $\int_S da |a|^{-2}$ can be for a domain S with diameter $2|w|$ and which is bounded away from the origin by $|w|^{\frac{1}{2}}$. Finally, since $|w| - |k| \leq |w|$ and by (3.10)

$$\left| \int_{-w}^w da \frac{k + a - S(k + a) (|k + a|^2 + 1)^{\frac{1}{2}}}{(|k + a|^2 + 1)^{\frac{1}{2}}} \right| \leq |2(|w| - |k|) + O(|w|^{-\frac{1}{2}})|^2 < 5|w| N(w, k).$$

For k such that $|w| - |k| \approx 0$, $O(|w|^{-\frac{1}{2}})$ may become the larger term, but it is still smaller than $N(w, k)$ which will be ≈ 1 when $|k|$ is near $|w|$. $\square$

We now apply Lemma 3.2 to attain a similar inequality with $|M'_s|$ replaced by $E_s^{\frac{1}{2}}$. Our analysis is based on the fact that $E_t^{\frac{1}{2}} = (\frac{1}{2}K_t^2 + V(X_t))^{\frac{1}{2}}$ is a submartingale. The submartingale property of $E_t^{\frac{1}{2}}$ follows since $f(k) = (\frac{1}{2}k^2 + V)^{\frac{1}{2}}$ is a convex function. Since the jumps occurs with symmetric probabilities $\mathcal{P}_a(v) = \mathcal{P}_a(-v)$, we can write $E_t^{\frac{1}{2}} - E_0^{\frac{1}{2}} = \mathbf{M}_t + \mathbf{A}_t$ in terms of a martingale part $\mathbf{M}_t$ and a stochastically increasing part $\mathbf{A}_t$ as

$$\mathbf{M}_t = \frac{1}{2} \sum_{n=1}^{\mathcal{N}_t} \left[\left(\frac{1}{2} |K_{t_n^-} + w_n|^2 + V(X_{t_n}) \right)^{\frac{1}{2}} - \left(\frac{1}{2} |K_{t_n^-} - w_n|^2 + V(X_{t_n}) \right)^{\frac{1}{2}} \right], \quad (3.13)$$

$$\begin{aligned} \mathbf{A}_t = \frac{1}{2} \sum_{n=1}^{\mathcal{N}_t} & \left[\left(\frac{1}{2} |K_{t_n^-} + w_n|^2 + V(X_{t_n}) \right)^{\frac{1}{2}} + \left(\frac{1}{2} |K_{t_n^-} - w_n|^2 + V(X_{t_n}) \right)^{\frac{1}{2}} \right. \\ & \left. - 2 \left(\frac{1}{2} |K_{t_n^-}|^2 + V(X_{t_n}) \right)^{\frac{1}{2}} \right], \end{aligned} \quad (3.14)$$

where t_n, w_n for $n = 1, \dots, \mathcal{N}_t$ are the Poisson times and their corresponding momentum jumps (when they occur), respectively. The processes $\mathbf{M}_t, \mathbf{A}_t$ form a Doob-Meyer decomposition for $E_t^{\frac{1}{2}} - E_0^{\frac{1}{2}}$, however $\mathbf{A}_t$ is not a predictable process so the decomposition is not in the unique sense.

Lemma 3.4 (Energy Lemma). *Assume List 2.1. Define $T_{\epsilon, V}^{(t)} = \int_0^1 ds \chi(|t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}}| > \epsilon)$. There exists a constant C such that*

$$\liminf_{t \rightarrow \infty} \Pr[T_{\epsilon, V}^{(t)} \geq 1 - \delta] \geq 1 - C \frac{r_2^{\frac{1}{2}}}{r_1} \frac{\epsilon}{\delta}.$$

Proof. Define the martingale M'_t and the increasing process A'_t as

$$M'_t = \sum_{n=1}^{\mathcal{N}_t} w_n S(K_{t_n^-}) \text{ and } A'_t = \sup_{0 \leq s \leq t} -M'_s,$$

where $S(K_{t-})$ is the left limit up to time t for the sign of the momentum. Also define $G_s^{(t)}$ as the difference $G_s^{(t)} = \sqrt{2}t^{-\frac{1}{2}}E_{st} - t^{-\frac{1}{2}}M'_{st} - t^{-\frac{1}{2}}A'_{st}$. In general, we have that

$$\Pr[T_{\epsilon}^{(t)} > 1 - \delta, \sup_{0 \leq s \leq 1} G_s^{(t)} \leq (\sqrt{2} - 1)\epsilon] \leq \Pr[T_{\epsilon, V}^{(t)} > 1 - \delta], \quad (3.15)$$

where $T_{\epsilon}^{(t)}$ is defined as in Lemma 3.2 for the process $t^{-\frac{1}{2}}M'_{st} + t^{-\frac{1}{2}}A'_{st}$. We will prove below that $\Pr[\sup_{0 \leq s \leq 1} G_s^{(t)} > (\sqrt{2} - 1)\epsilon] = O(t^{-\frac{1}{4}})$. In that case, by applying the inclusion-exclusion principle to the left side of (3.15), then (3.15) can be written

$$\Pr[T_{\epsilon}^{(t)} > 1 - \delta] + O(t^{-\frac{1}{4}}) \leq \Pr[T_{\epsilon, V}^{(t)} > 1 - \delta].$$

We can then apply Lemma 3.2 to the left-side to complete the proof.

Now we work towards establishing $\Pr [\sup_{0 \leq s \leq 1} G_s^{(t)} > (\sqrt{2} - 1)\epsilon] = O(t^{-\frac{1}{4}})$. Consider the martingale $M'_t - \mathbf{M}_t$. The square of the jumps of $M'_t - \mathbf{M}_t$ can be bounded by the jumps of $\mathbf{A}_t$ plus an extra term through the inequality from Lemma 3.3:

$$\begin{aligned} & \left| \sqrt{2} w S(k) - \left[\left(\frac{1}{2} |k + w|^2 + V(x) \right)^{\frac{1}{2}} - \left(\frac{1}{2} |k - w|^2 + V(x) \right)^{\frac{1}{2}} \right] \right|^2 \\ & \leq 2 v^2 1_{|w| > J} + c J \left[\left(\frac{1}{2} |k + w|^2 + V(x) \right)^{\frac{1}{2}} + \left(\frac{1}{2} |k - w|^2 + V(x) \right)^{\frac{1}{2}} - 2 \left(\frac{1}{2} k^2 + V(x) \right)^{\frac{1}{2}} \right]. \end{aligned} \quad (3.16)$$

for all $J > 1$, $k, w, x \in \mathbb{R}$. Define the process $Q_r = \sum_{n=1}^{\mathcal{N}_r} w_n^2 \chi(|w_n| \geq t^{\frac{1}{4}})$.

By (3.16), the quadratic variation process $[M' - \mathbf{M}]_t$ has the bound

$$[M' - \mathbf{M}]_t \leq c t^{\frac{1}{4}} \mathbf{A}_t + 2Q_r.$$

This will allow us to bound $\mathbb{E}[(M'_t - \mathbf{M}_t)^2] = \mathbb{E}[M' - \mathbf{M}]_t$. By the fact that $\mathbf{A}_t$ is the increasing part of the Doob-Meyer decomposition for $E_t^{\frac{1}{2}} - E_0^{\frac{1}{2}}$ and Jensen's inequality we have

$$\mathbb{E}[\mathbf{A}_t] = \mathbb{E}[E_t^{\frac{1}{2}} - E_0^{\frac{1}{2}}] \leq (\mathbb{E}[E_t])^{\frac{1}{2}} \leq (\mathbb{E}[E_0] + 2^{-1} r_2 t)^{\frac{1}{2}} \sim 2^{-\frac{1}{2}} r_2^{\frac{1}{2}} t^{\frac{1}{2}}.$$

Also we have

$$\mathbb{E}[Q_t] = \mathbb{E}\left[\sum_{n=1}^{\mathcal{N}_t} w_n^2 \chi(|w_n| \geq t^{\frac{1}{4}})\right] \leq \mathbb{E}[\mathcal{N}_t] \rho t^{-\frac{1}{2}} = \mathcal{R} \rho t^{\frac{1}{2}},$$

where we have used that the fourth moments of the jumps are bounded as $\mathbb{E}[w_n^4] \leq \rho$, since

$$\mathbb{E}[w_n^2 \chi(|w_n| \geq t^{\frac{1}{4}})] \leq t^{-\frac{1}{2}} \mathbb{E}[w_n^4 \chi(|w_n| \geq t^{\frac{1}{4}})] \leq t^{-\frac{1}{2}} \rho.$$

Thus, using the above and Doob's maximal inequality

$$\begin{aligned} \mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-1} |M'_{st} - \mathbf{M}_{st}|^2\right] & \leq 4t^{-1} \mathbb{E}[(M'_t - \mathbf{M}_t)^2] = 4\mathbb{E}[M' - \mathbf{M}]_t \\ & \leq 4c t^{-\frac{3}{4}} \mathbb{E}[\mathbf{A}_t] + 8t^{-1} \mathbb{E}[Q_t] \leq 4c 2^{-\frac{1}{2}} r_2^{\frac{1}{2}} t^{-\frac{1}{4}} + 8\mathcal{R} \rho t^{-\frac{1}{2}} = O(t^{-\frac{1}{4}}). \end{aligned} \quad (3.17)$$

Next we show that $t^{-\frac{1}{2}} \mathbf{A}_{st}$ is typically bounded from below by $t^{-\frac{1}{2}} A'_{st}$ in the sense

$$\mathbb{E}\left[\sup_{0 \leq s \leq 1} |(t^{-\frac{1}{2}} A'_{st} - t^{-\frac{1}{2}} \mathbf{A}_{st}) 1_{A'_{st} > \mathbf{A}_{st}}|^2\right] = O(t^{-\frac{1}{4}}).$$

We can write $\sqrt{2} t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}}$ as

$$\sqrt{2} t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}} = t^{-\frac{1}{2}} M'_{st} + \sqrt{2} t^{-\frac{1}{2}} E_0^{\frac{1}{2}} + t^{-\frac{1}{2}} \mathbf{M}_{st} - t^{-\frac{1}{2}} M'_{st} + t^{-\frac{1}{2}} \mathbf{A}_{st} \quad (3.18)$$

where since the left side is positive for all a , we must have

$$-t^{-\frac{1}{2}} M'_{at} \chi(t^{-\frac{1}{2}} M'_{at} \leq 0) \leq (\sqrt{2} t^{-\frac{1}{2}} E_0^{\frac{1}{2}} + t^{-\frac{1}{2}} \mathbf{M}_{at} - t^{-\frac{1}{2}} M'_{at} + t^{-\frac{1}{2}} \mathbf{A}_{at}) \chi(t^{-\frac{1}{2}} M'_{at} \leq 0).$$

Taking the supremum in a up to $s \leq 1$ of both sides,

$$t^{-\frac{1}{2}} A'_{st} = \sup_{0 \leq a \leq s} -t^{-\frac{1}{2}} M'_{at} \leq \sup_{0 \leq a \leq s} |\sqrt{2} t^{-\frac{1}{2}} E_0^{\frac{1}{2}} + t^{-\frac{1}{2}} \mathbf{M}_{at} - t^{-\frac{1}{2}} M'_{at}| + t^{-\frac{1}{2}} \mathbf{A}_{st},$$

where we have used that $M'_0 = 0$ for the left side and that $\mathbf{A}_r$ is increasing for the right side. Subtracting $t^{-\frac{1}{2}}\mathbf{A}_{st}$ and taking $\mathbb{E}[\sup_{0 \leq s \leq 1} |\cdot|^2]^{\frac{1}{2}}$ of both sides

$$\begin{aligned} \mathbb{E}\left[\sup_{0 \leq s \leq 1} \left|(t^{-\frac{1}{2}}A'_{st} - t^{-\frac{1}{2}}\mathbf{A}_{st})1_{A'_{st} > \mathbf{A}_{st}}\right|^2\right]^{\frac{1}{2}} \\ \leq \sqrt{2}t^{-\frac{1}{2}}E_0^{\frac{1}{2}} + \mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-1}|\mathbf{M}_{st} - M'_{st}|^2\right]^{\frac{1}{2}} = O(t^{-\frac{1}{8}}). \end{aligned} \quad (3.19)$$

Observe that $G_s^{(t)} < 0$ implies

$$\mathbf{A}_{st} - A'_{st} \leq \sqrt{2}E_0^{\frac{1}{2}} - 0 \wedge (\mathbf{M}_{st} - M'_{st}). \quad (3.20)$$

Finally, by the triangle inequality, (3.17), (3.19), and (3.20)

$$\begin{aligned} \mathbb{E}\left[\sup_{0 \leq s \leq 1} |G_s^{(t)}|^2 1_{G_s^{(t)} < 0}\right]^{\frac{1}{2}} \\ \leq t^{-\frac{1}{2}}\sqrt{2}E_0^{\frac{1}{2}} + \mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-1}|\mathbf{A}_{st} - A'_{st}|^2 1_{G_s^{(t)} < 0}\right]^{\frac{1}{2}} + \mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-1}|\mathbf{M}_{st} - M'_{st}|^2\right]^{\frac{1}{2}} \\ \leq t^{-\frac{1}{2}}\sqrt{2}E_0^{\frac{1}{2}} + \mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-\frac{1}{2}}|\mathbf{A}_{st} - A'_{st}|^2 1_{A'_{st} > \mathbf{A}_{st}}\right]^{\frac{1}{2}} + 2\mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-1}|\mathbf{M}_{st} - M'_{st}|^2\right]^{\frac{1}{2}}. \end{aligned} \quad (3.21)$$

The right-side is $O(t^{-\frac{1}{8}})$, and via Chebyshev's inequality $\Pr[\sup_{0 \leq s \leq 1} |G_s^{(t)}| > (\sqrt{2} - 1)\epsilon] = O(t^{-\frac{1}{4}})$, which is the bound that was claimed. $\square$

In the following lemma, we set $L_{st} = M'_{st} + A'_{st}$ where M'_{st} and A'_{st} are defined as in the proof of Lemma 3.4. The Doob-Meyer decompositions in the lemma are not unique, since the increasing parts are not constrained to be predictable.

Lemma 3.5. *Consider the submartingales E_t and L_t^2 .*

1. E_t admits a Doob-Meyer decomposition as a sum of martingale and increasing parts

$$\mathcal{M}_t = \sum_{n=1}^{\mathcal{N}_t} w_n K_{t_n^-} \quad \text{and} \quad \mathcal{A}_t = E_0 + \frac{1}{2} \sum_{n=1}^{\mathcal{N}_t} w_n^2.$$

2. L_t^2 admits a Doob-Meyer decomposition as a sum of martingale and increasing parts $\bar{M}_t = \sum_{n=1}^{\mathcal{N}_t} \alpha_n$ and $\bar{A}_t = \sum_{n=1}^{\mathcal{N}_t} \beta_n$ respectively, where

$$\begin{aligned} \alpha_n &= 2w_n L_{t_{n-1}} & \text{for } |L_{t_{n-1}}| \geq |w_n|, \\ \alpha_n &= \frac{1}{2} S(K_{t_n^-}) S(w_n) (L_{t_{n-1}} + |w_n|)^2 & \text{for } |L_{t_{n-1}}| \leq |w_n|, \\ \beta_n &= w_n^2 & \text{for } |L_{t_{n-1}}| \geq |w_n|, \\ \beta_n &= w_n^2 + \frac{1}{2} (|w_n| - L_{t_{n-1}})^2 & \text{for } |L_{t_{n-1}}| \leq |w_n|. \end{aligned}$$

3. In the limit $t \rightarrow \infty$,

$$\mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-1} |2\mathcal{A}_{st} - \bar{A}_{st}|\right] = O(t^{-\frac{1}{4}}).$$

Proof.

Part (1): E_t can be rewritten as

$$E_t = E_0 + \sum_{n=1}^{\mathcal{N}_t} w_n K_{t_n^-} + \frac{1}{2} w_n^2, \quad (3.22)$$

which follows by an inductive expansion using the conservation of energy between momentum jumps:

$$\begin{aligned} E_{t_n} = \frac{1}{2}(K_{t_n^-} + w_n)^2 + V(X_{t_n^-}) &= \frac{1}{2}K_{t_n^-}^2 + V(X_{t_n^-}) + w_n K_{t_n^-} + \frac{1}{2}w_n^2 \\ &= \frac{1}{2}K_{t_{n-1}}^2 + V(X_{t_{n-1}}) + w_n K_{t_n^-} + \frac{1}{2}w_n^2. \end{aligned}$$

The middle term on the right of (3.22) is a martingale by the symmetry of the jump rates (III) of List 2.1: $E[w_n | \mathcal{F}_{t_n^-}] = 0$.

Part (2): To find an expression for L_t , we apply an inductive argument as with E_t except that the analysis breaks into two cases. Expanding $L_{t_n}^2$ when $|L_{t_{n-1}}| > |w_n|$ is the easy case since $L_{t_n}^2 = L_{t_{n-1}}^2 + 2w_n S(K_{t_n^-}) + w_n^2$. Again $w_n S(K_{t_n^-})$ is the martingale contribution, since $E[w_n^2 | \mathcal{F}_{t_n^-}] = 0$.

Expanding $L_{t_n}^2$ in the case that $|L_{t_{n-1}}| \leq |w_n|$, then $L_{t_n} = \frac{1}{2}(1 + S(K_{t_n^-})S(w_n))(L_{t_{n-1}} + |w_n|)$, and

$$L_{t_n}^2 - L_{t_{n-1}}^2 = \frac{1}{2}S(K_{t_n^-})S(w_n)(L_{t_{n-1}} + |w_n|)^2 + w_n^2 + \frac{1}{2}(|w_n| - L_{t_{n-1}})^2.$$

The first term on the right has mean zero since $\mathbb{E}[S(w_n) | \mathcal{F}_{t_n^-}, |w_n|] = 0$.

Part (3): By Part (1) and Part (2),

$$t^{-1}(\bar{\mathcal{A}}_{st} - 2\mathcal{A}_{st}) = -2t^{-1}E_0 + t^{-1} \sum_{n=1}^{\mathcal{N}_{st}} \frac{1}{2}(|w_n| - L_{t_{n-1}})^2 \chi(|w_n| \geq L_{t_{n-1}})$$

Since $|w_n| \leq J$, the sum above is bounded by

$$\begin{aligned} &\sum_{n=1}^{\mathcal{N}_{st}} (|w_n| - L_{t_{n-1}})^2 \chi(|L_{t_{n-1}}| \leq |w_n|) \\ &\leq t^{\frac{1}{4}} \sum_{n=1}^{\mathcal{N}_{st}} (|w_n| - L_{t_{n-1}}) \chi(|L_{t_{n-1}}| \leq |w_n|) + \sum_{n=1}^{\mathcal{N}_{st}} |w_n|^2 \chi(|w_n| \geq t^{\frac{1}{4}}). \quad (3.23) \end{aligned}$$

By the estimates in Lemma 3.4

$$\mathbb{E} \left[\sup_{0 \leq s \leq 1} t^{-1} \sum_{n=1}^{\mathcal{N}_{st}} |w_n|^2 \chi(|w_n| \geq t^{\frac{1}{4}}) \right] = \mathbb{E} \left[t^{-1} \sum_{n=1}^{\mathcal{N}_t} |w_n|^2 \chi(|w_n| \geq t^{\frac{1}{4}}) \right] = O(t^{-\frac{1}{2}}).$$

The first term on the right-side is closely related to A'_r , since A'_r can be written

$$\sup_{0 \leq s \leq r} - \sum_{n=1}^{\mathcal{N}_r} w_n S(K_{t_n^-}) = A'_r = \sum_{n=1}^{\mathcal{N}_r} (|w_n| - L_{t_{n-1}}) \chi(|L_{t_{n-1}}| \leq |w_n|, S(K_{t_n^-}) S(w_n) = -1),$$

since increases in A'_t occur $|L_{t_{n-1}}| \leq |w_n|$ and the jump w_n has sign such that $S(K_{t_n^-}) S(w_n) = -1$. In fact, the conditional expectation of a single term from the sum with respect to the information up to time t_n^- and the size $|w_n|$ of the n th jump is

$$\begin{aligned} \mathbb{E}[(|w_n| - L_{t_{n-1}}) \chi(|L_{t_{n-1}}| \leq |w_n|, S(K_{t_n^-}) S(w_n) = -1) \mid \mathcal{F}_{t_n^-}, |w_n|] \\ = \frac{1}{2}(|w_n| - L_{t_{n-1}}) \chi(|L_{t_{n-1}}| \leq |w_n|), \end{aligned}$$

since $w_n = -|w_n|$ and $w_n = |w_n|$ have equal probability. Thus,

$$\begin{aligned} \mathbb{E}\left[t^{-\frac{3}{4}} \sum_{n=1}^{\mathcal{N}_t} (|w_n| - L_{t_{n-1}}) \chi(|L_{t_{n-1}}| \leq |w_n|)\right] &= t^{-\frac{3}{4}} \mathbb{E}[A'_t] = t^{-\frac{3}{4}} \mathbb{E}[M'_t + A'_t] \\ &= \mathbb{E}\left[2^{\frac{1}{2}} t^{-\frac{3}{4}} E_t^{\frac{1}{2}} - t^{-\frac{1}{4}} G_1^{(t)}\right] \leq 2^{\frac{1}{2}} t^{-\frac{3}{4}} \mathbb{E}[E_t^{\frac{1}{2}}] + O(t^{-\frac{3}{8}}) \\ &\leq 2^{\frac{1}{2}} t^{-\frac{3}{4}} \mathbb{E}[E_t]^{\frac{1}{2}} + O(t^{-\frac{3}{8}}) \leq 2^{\frac{1}{2}} t^{-\frac{3}{4}} (\mathbb{E}[E_0] + \frac{1}{2} r_2 t)^{\frac{1}{2}} + O(t^{-\frac{3}{8}}) = O(t^{-\frac{1}{4}}), \quad (3.24) \end{aligned}$$

where the second equality is because M'_r is a mean zero martingale, the third equality is from the definition of $G_r^{(t)}$, and the first inequality uses the result $\mathbb{E}[\sup_{0 \leq s \leq 1} |G_s^{(t)}|^2 1_{G_s^{(t)} < 0}]^{\frac{1}{2}} = O(t^{-\frac{1}{8}})$ from the proof of Lemma 3.4. $\square$

4 The absolute value of the momentum

The following theorem takes us as far possible towards attaining a central limit theorem for the momentum $t^{-\frac{1}{2}} K_{st}$ without making an assumption about a reflection symmetry in the periodic potential and in the jump probabilities.

Theorem 4.1. *Assume List 2.1 and (i)-(ii) of List 2.2. In the limit $t \rightarrow \infty$, $t^{-\frac{1}{2}} |K_{st}|$ converges in distribution to the absolute value of a Brownian motion with diffusion constant $\sigma = \int_{\mathbb{R}} dv \tilde{P}(v) v^2$.*

The above theorem is not enough to guarantee that $t^{-\frac{1}{2}} K_{st}$ converges to a Brownian motion. To see this, consider a random walk X_n on $\mathbb{Z}$ which jumps to the right and the left with equal probability at every lattice site except at the origin where it jumps to 1 with probability $\frac{2}{3}$ and to -1 otherwise. In this case, $|X_n|$ has the same distribution as a simple random walk and thus $n^{-\frac{1}{2}} |X_{[sn]}|$ converges in distribution to the absolute value of a Brownian motion $|\mathbf{B}_s|$. However, removing the absolute values, then $n^{-\frac{1}{2}} X_{[sn]}$ will have a drift determined by the process $\frac{1}{3} n^{-\frac{1}{2}} \sum_{n=1}^{[sn]} \chi(X_n = 0)$. Since a simple random walk X_n spends on the order of $n^{\frac{1}{2}}$ steps at the origin, the drift will be non-vanishing.

The proof of Theorem 4.1 follows in Section 7 and is based on the Lemma 4.2 below. The latter extends the analysis in the proof of Lemma 3.4 to show that $t^{-\frac{1}{2}} K_{st}$ is close to being

the solution of a stochastic equation, reminiscent of the fact for a Brownian motion $\mathbf{B}_s$ that $|\mathbf{B}_s| = \mathbf{B}'_s + \sup_{0 \leq r \leq s} -\mathbf{B}'_s$ for another Brownian motion $\mathbf{B}'_s = \int_0^s S(\mathbf{B}_r) d\mathbf{B}_r$. This is related to Levy's theory of Brown local time [11].

For some bounded right-continuous adapted process F_t , we denote

$$\int_0^s F_{rt} dM_r^{(t)} = t^{-\frac{1}{2}} \sum_{n=1}^{\mathcal{N}_{st}} w_n F_{t_n}^-, \quad (4.1)$$

where v_n are the momentum jumps. Thus the martingale M'_t used in the proof of Lemma 3.4 can be written $t^{-\frac{1}{2}} M'_{st} = \int_0^s S(K_{rt}) dM_r^{(t)}$.

Lemma 4.2. *Assume List 2.1. Then the momentum process K_{st} , $s \in [0, 1]$, satisfies the stochastic equation*

$$t^{-\frac{1}{2}} |K_{st}| = \int_0^s S(K_{rt}) dM_r^{(t)} + \sup_{0 \leq a \leq s} - \int_0^a S(K_{rt}) dM_r^{(t)} + \mathcal{E}_s^{(t)}, \quad (4.2)$$

where the error $\mathcal{E}_s^{(t)}$ has $\mathbb{E}[\sup_{0 \leq s \leq 1} |\mathcal{E}_s^{(t)}|^2] \rightarrow 0$ for large t .

Proof. The error $\mathcal{E}_s^{(t)}$ is close to error $G_s^{(t)} = \sqrt{2} t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}} - t^{-\frac{1}{2}} M'_{st} - t^{-\frac{1}{2}} A'_{st}$ which arose in the proof of Lemma 3.4, since they differ only by the replacement of the momentum $t^{-\frac{1}{2}} |K_{st}|$ with $\sqrt{2} t^{\frac{1}{2}} E_{st}^{\frac{1}{2}}$ and $|\sqrt{2} t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}} - t^{-\frac{1}{2}} |K_{st}|| \leq t^{-\frac{1}{2}} \bar{V}$. In the proof of Lemma 3.4, it was shown that

$$t^{-1} \mathbb{E} \left[\sup_{0 \leq s \leq 1} |G_s^{(t)}|^2 1_{G_s^{(t)} < 0} \right] = O(t^{-\frac{1}{4}}). \quad (4.3)$$

We now work on showing that $t^{-1} \mathbb{E}[\sup_{0 \leq s \leq 1} |G_s^{(t)}|^2 1_{G_s^{(t)} \geq 0}]$ tends to zero also.

By Part (3) of Lemma 3.5, the normalized difference in the increasing parts $\bar{\mathcal{A}}_{st}$ and $\mathcal{A}_{st}$ for the Doob-Meyer decompositions of L_{st}^2 and E_{st} respectively, tends to zero as

$$\mathbb{E} \left[\sup_{0 \leq s \leq 1} t^{-1} |2\mathcal{A}_{st} - \bar{\mathcal{A}}_{st}| \right] = O(t^{-\frac{1}{4}}). \quad (4.4)$$

We will make use of the error $G_s^{(t)}$ being the difference between two positive submartingales having a vanishing difference between the increasing parts of their Doob-Meyer decompositions:

$$\begin{aligned} H_s^{(t)} &= 2t^{-1} E_{st} - t^{-1} |M'_{st} + A'_{st}|^2 = 2t^{-1} E_{st} - (\sqrt{2} t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}} - G_s^{(t)})^2 \\ &= G_s^{(t)} (2^{\frac{3}{2}} t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}} - G_s^{(t)}) = \Upsilon_s^{(t)} + \Gamma_s^{(t)}, \end{aligned}$$

where $\Upsilon_s^{(t)}$ is a martingale and $\Gamma_s^{(t)} = t^{-1} (2\mathcal{A}_{st} - \bar{\mathcal{A}}_{st})$ is $O(t^{-\frac{1}{4}})$ by (4.4).

Note that $2^{\frac{3}{2}} t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}} - G_s^{(t)}$ is positive, so $G_s^{(t)}$ determines the sign of $H_s^{(t)}$. By the Cauchy-Schwarz inequality

$$t^{-1} \mathbb{E} \left[\sup_{0 \leq s \leq 1} |H_s^{(t)}| 1_{H_s^{(t)} < 0} \right] \leq \left(\mathbb{E} \left[\sup_{0 \leq s \leq 1} |G_s^{(t)}|^2 1_{G_s^{(t)} < 0} \right] \right)^{\frac{1}{2}} \left(\mathbb{E} \left[\sup_{0 \leq s \leq 1} |2^{\frac{3}{2}} t^{-\frac{1}{2}} E_{st}^{\frac{1}{2}} - G_s^{(t)}|^2 \right] \right)^{\frac{1}{2}}. \quad (4.5)$$

The first factor on the right is $O(t^{-\frac{1}{8}})$ by (4.3). Bounding the second factor on the right comes through the triangle inequality and the use of Doob's maximal inequality for the two positive submartingales $M_{st} + A_{st}$ and $E_{st}^{\frac{1}{2}}$:

$$\left(\mathbb{E}\left[\sup_{0 \leq s \leq 1} \left|2^{\frac{3}{2}}t^{-\frac{1}{2}}E_{st}^{\frac{1}{2}} - G_s^{(t)}\right|^2\right]\right)^{\frac{1}{2}} \leq 4\sqrt{2}t^{-\frac{1}{2}}\mathbb{E}[E_{st}]^{\frac{1}{2}} + 4t^{-\frac{1}{2}}\mathbb{E}[|M'_{st} + A'_{st}|^2]^{\frac{1}{2}} \leq 12r_2^{\frac{1}{2}}, \quad (4.6)$$

where $2^{\frac{3}{2}}t^{-\frac{1}{2}}E_{st}^{\frac{1}{2}} - G_s^{(t)} = 2^{\frac{1}{2}}t^{-\frac{1}{2}}E_{st}^{\frac{1}{2}} + t^{-\frac{1}{2}}M'_{st} + t^{-\frac{1}{2}}A'_{st}$. The last inequality follows by the forms for the increasing parts of the Doob-Meyer decompositions for $t^{-1}|M'_{st} + A'_{st}|^2$ and $t^{-1}E_{st}$ from Lemma 3.5 which are both bounded by $t^{-1}[M]_{st}$. The right side of (4.5) vanishes as $O(t^{-\frac{1}{8}})$, and thus the values of $t^{-1}H_s^{(t)}$ do not typically go far in the negative direction. Since $H_s^{(t)}$ is a martingale with mean $2t^{-1}E_0 \rightarrow 0$, it would be expected that $t^{-1}H_s^{(t)}$ can also not go far in the positive direction.

We will argue below that $\lim_{t \rightarrow \infty} \mathbb{E}[\sup_{0 \leq s \leq 1} |H_s^{(t)}| 1_{H_s^{(t)} > 0}] \rightarrow 0$. This would complete the proof since

$$t^{-1}\mathbb{E}\left[\sup_{0 \leq s \leq 1} |G_s^{(t)}|^2 1_{G_{st} > 0}\right] \leq t^{-1}\mathbb{E}\left[\sup_{0 \leq s \leq 1} H_s^{(t)} 1_{H_s^{(t)} > 0}\right],$$

which clearly follows since $2^{\frac{3}{2}}t^{-\frac{1}{2}}E_{st}^{\frac{1}{2}} - G_s^{(t)}$ is larger in absolute value than $G_s^{(t)}$ when $G_s^{(t)}$ is positive.

By the optional sampling theorem,

$$\mathbb{E}[\Upsilon_\tau^{(t)}] = \mathbb{E}[H_\tau^{(t)} - \Gamma_\tau^{(t)}] = 0 \quad (4.7)$$

for any adapted stopping time τ . Given some $a > 0$, let $\tau \in [0, 1]$ be the first time that $H_s^{(t)}$ reaches above a and put $\tau = 1$ if that event does not occur. By (4.7),

$$\mathbb{E}[\Gamma_\tau^{(t)}] = a \Pr\left[\sup_{0 \leq s \leq 1} H_s^{(t)} \geq a\right] + \mathbb{E}[H_1^{(t)} \chi(\sup_{0 \leq s \leq 1} H_s^{(t)} < a)],$$

which implies that

$$\sup_{a \in \mathbb{R}^+} a \Pr\left[\sup_{0 \leq s \leq 1} H_s^{(t)} \geq a\right] \leq |\mathbb{E}[\Gamma_\tau^{(t)}]| + \mathbb{E}\left[\sup_{0 \leq s \leq 1} |H_s^{(t)}| 1_{H_s^{(t)} < 0}\right] = O(t^{-\frac{1}{4}}). \quad (4.8)$$

Set $Y_t = \sup_{0 \leq s \leq 1} H_s^{(t)} 1_{H_s^{(t)} > 0}$ and $\alpha_t = (\sup_{a \in \mathbb{R}^+} a \Pr[Y_t \geq a])^{\frac{1}{2}}$. If $\mathbb{E}[Y_t^2] \leq \mathbf{y}^2$ for all t , then for any $m > \alpha_t > 0$

$$\begin{aligned} \mathbb{E}[Y_t] &= \int_0^\infty da \Pr[Y_t \geq a] = \left(\int_0^{\alpha_t} + \int_{\alpha_t}^m + \int_m^\infty\right) da \Pr[Y_t \geq a] \\ &\leq \alpha_t + \frac{1}{\alpha_t} \int_{\alpha_t}^m da a \Pr[Y_t \geq a] + \frac{1}{m} \int_m^\infty da a \Pr[Y_t \geq a] \leq \alpha_t + m\alpha_t + \frac{\mathbf{y}^2}{2m}. \end{aligned}$$

The second inequality above uses that α_t^2 is the supremum of $a \Pr[Y_t \geq a]$ for the first integral and the relation $\frac{1}{2}\mathbb{E}[Y_t^2] = \int_0^\infty da a \Pr[Y_t \geq a]$ for the second. We can pick m large to make the last term on the right-side small, and then pick t large enough so that with (4.8) $\alpha_t(1+m)$ is small.

To finish the argument we just need to show that $\mathbb{E}[Y_t^2]$ can be uniformly bounded in t . Since $G_\tau^{(t)}$ is the difference of $2^{\frac{1}{2}}t^{-\frac{1}{2}}E_{st}^{\frac{1}{2}}$ and $t^{-\frac{1}{2}}L_{st}$, and $2^{\frac{3}{2}}t^{-\frac{1}{2}}E_{st}^{\frac{1}{2}} - G_s^{(t)}$ is their sum,

$$\mathbb{E}[Y_t^2]^{\frac{1}{2}} < 4\mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-2}E_{st}^2\right]^{\frac{1}{2}} + 2\mathbb{E}\left[\sup_{0 \leq s \leq 1} t^{-2}|L_{st}|^4\right]^{\frac{1}{2}}.$$

To attain a value $\mathbf{y}$, the terms on the right can be bounded by standard calculations using the Doob-Meyer decompositions for E_r and $|L_r|^2$ from Lemma 3.5 and using Doob's maximal inequality.

□

5 Estimates at high energy

In this section, we provide estimates that are useful for understanding the dynamics when the particle has high energy, which by Lemma 3.4, is the majority of time. The estimates are based on the idea that the particle will feel a spatially averaged noise and that the momentum is too high to be shifted through the action of the force generated by the potential.

The following elementary bound is used many times in this section and later sections. It follows from the conservation of energy and the quadratic formula. It basically says that if the initial momentum k_0 has $|k_0| \gg \bar{V}^{\frac{1}{2}} = \sup_{a \in [0,1]} V(x)^{\frac{1}{2}}$, then the future momenta k_s , as determined by the Hamiltonian evolution, will stay close to k_0 .

Lemma 5.1. *Let (x_t, k_t) evolve according to the Hamiltonian $H(x, k) = \frac{1}{2}k^2 + V(x)$, for positive potential bounded by $\bar{V}$. If the initial momentum has $|k_0|^2 > 4\bar{V}$, then the displacements in momentum $k_t - k_0$ and $k_t - k_s$ have bounds*

$$|k_t - k_0| < \frac{2\bar{V}}{|k_0|} \text{ and } |k_t - k_s| < \frac{4\bar{V}}{|k_0|} \text{ for all } t, s \in \mathbb{R}.$$

Proof. Since $|k_0|^2 > 4\bar{V}$, the momentum k_t will not change signs at any time. By the conservation of energy

$$\frac{1}{2}|k_0 + (k_t - k_0)|^2 - \frac{1}{2}k_0^2 = -V(x_t) + V(x_0).$$

Using the quadratic formula and that k_t, k_0 have the same sign,

$$|k_t - k_0| = \left| |k_0| - (k_0^2 + 2V(x_0) - 2V(x_t))^{\frac{1}{2}} \right| \leq \left| \frac{1}{2} \int_0^{2V(x_0) - 2V(x_t)} da (k_0^2 + a)^{-\frac{1}{2}} \right| < \frac{2\bar{V}}{|k_0|}$$

Since $(\frac{1}{2}k_0^2 + a)^{-\frac{1}{2}} \leq \sqrt{2}|k_0|^{-1} < 2|k_0|^{-1}$ for $a \leq \frac{1}{2}k_0^2$. By the triangle inequality, we can bound the difference $|k_t - k_s|$.

□

As before $\tilde{P}(v) = \int_0^1 da \frac{\kappa(a)}{\bar{\kappa}} \mathcal{P}_a(v)$ where $\bar{\kappa} = \int_0^1 da \kappa(a)$. By our assumptions $\inf_{0 \leq a \leq 1} \kappa(a) = \nu > 0$.

Lemma 5.2. *Assume List 2.1 and (i)-(ii) of List 2.2. Starting from the point (x, k) with $|k|^2 \gg \bar{V}$, let $r_{(x,k)} \in L^1([0, 1])$ and $\tilde{r}_{(x,k)} \in L^1([0, 1])$ be the probability density for the torus position of the particle at the first Poisson time and at the time of the first momentum jump respectively. Let $P_{(x,k)} \in L^1(\mathbb{R})$ be the density for the first momentum jump. Let $T_{(x,k),t} \in L^1([0, 1])$ be the density of time that a deterministic trajectory starting from (x, k) and evolving according to the Hamiltonian $H(x, k) = k^2 + V(x)$ spends at a torus point over a time interval $[0, t]$. We have the following bounds*

$$1. \sup_{a \in [0, 1]} |r_{(x,k)}(a) - 1| \leq \frac{2\mathcal{R}}{|k|} + O\left(\frac{1}{|k|^2}\right),$$

2. $\sup_{a \in [0, 1]} \left| \tilde{r}_{(x, k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \leq \frac{2\mathcal{R}\nu^{-2}}{|k|} + O\left(\frac{1}{|k|^2}\right),$
3. $\sup_{v \in \mathbb{R}} \left| \frac{P_{(x, k)}(v)}{\bar{P}(v)} - 1 \right| \leq \frac{2\mathcal{R}\nu^{-2}}{|k|} + O\left(\frac{1}{|k|^2}\right).$
4. $\sup_{a \in [0, 1]} \left| T_{(x, k), t}(a) - t \right| \leq \frac{2t}{|k|}.$

Proof.

Part (1):

Let $(x_s, k_s) \in \mathbb{R}^2$ be the position and momentum for a particle beginning at (x, k) and evolving over a time period s for Hamiltonian $H(x, k) = \frac{1}{2}k^2 + V(x)$. Notice that $r_{(x, k)}(a)$ can be written as

$$r_{(x, k)}(a) = \sum_{n=1}^{\infty} |\mathbf{k}_{(x, k)}(a)|^{-1} \mathcal{R} e^{-\mathcal{R} r_n(a)},$$

where $s = r_1(a), r_2(a), \dots$ are the periodic sequence of times for which $x_s \bmod(1) = a$, and $\mathbf{k}_{(x, k)}(a) = s(k)(H(x, k) - V(a))^{\frac{1}{2}}$ is their momentum at the point. These times will exist for every $a \in [0, 1]$ as long as $H(x, k) > \bar{V}$.

If $4\bar{V} \leq k^2$, then $|k_s - k| \leq 2\bar{V}|k|^{-1}$ by Lemma 5.1. Thus for large momentum $|k| \gg (\bar{V})^{\frac{1}{2}}$, k_s is nearly constant and the hit times $r_n(a)$ will be close to the sequence of times $s = s_n(a)$ at which $x + sk \bmod(1) = a$ for a time period at least on the order of $t^{\frac{1}{2}}$. The period τ such that $r_n(a) - r_{n-1}(a) = \tau$ should thus be close to $\frac{1}{|k|}$. When $|k|$ is large enough so that $|k_s - k| \leq 2\bar{V}|k|^{-1} < \frac{1}{2}|k|$, then clearly $\tau \leq \frac{2}{|k|}$, and

$$\left| \tau - \frac{1}{|k|} \right| \leq \frac{1}{|k|} \left| \int_0^{\tau} ds k - \int_0^{\frac{1}{|k|}} ds k \right| \leq \frac{1}{|k|} \left(\int_0^{\tau} ds |k_s - k| + \int_0^{\frac{1}{|k|}} ds |k_s - k| \right) < \frac{6\bar{V}}{|k|^3}. \quad (5.1)$$

The difference between the first crossing-times $|r_1(a) - s_1(a)|$ of the point a can be similarly bounded.

Using the triangle inequality

$$\begin{aligned} |r_{(x, k)}(a) - 1| &\leq \left| r_{(x, k)}(a) - \frac{1}{|k|} \sum_{n=1}^{\infty} \mathcal{R} e^{-\mathcal{R} r_n(a)} \right| \\ &\quad + \left| \frac{1}{|k|} \sum_{n=1}^{\infty} \mathcal{R} e^{-\mathcal{R} r_n(a)} - \frac{1}{|k|} \sum_{n=1}^{\infty} \mathcal{R} e^{-\mathcal{R} s_n(a)} \right| + \left| \frac{1}{|k|} \sum_{n=1}^{\infty} \mathcal{R} e^{-\mathcal{R} s_n(a)} - 1 \right| \\ &\leq \frac{2\mathcal{R}}{|k|} + O\left(\frac{1}{|k|^2}\right), \end{aligned} \quad (5.2)$$

where the last inequality follows by further computation using the inequalities above. For instance, we can bound the first term on the right as

$$\left| r_{(x, k)} - \frac{1}{|k|} \sum_{n=1}^{\infty} \mathcal{R} e^{-\mathcal{R} r_n(a)} \right| \leq \left| 1 - \frac{k}{\mathbf{k}_{(x, k)}(a)} \right| e^{-\mathcal{R} r_1(a)} \frac{1}{|k|\tau} \frac{\mathcal{R}\tau}{1 - e^{-\mathcal{R}\tau}} \leq \frac{2\bar{V}}{k^2}.$$

Part (2):

We will bound $\sup_{a \in [0, 1]} |\tilde{r}_{(x, k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1|$ by invoking Part (1). This will involve breaking down an expression for $\tilde{r}_{(x, k)}(a)$. Since there are a random number of Poisson times before the time of the first momentum jump, the expression will have a series of integrals whose n th term corresponds to a momentum jump occurs at the $(n + 1)$ th Poisson time.

$$\tilde{r}_{(x, k)}(a) = \kappa(a) \sum_{n=0}^{\infty} \int_{(\mathbb{R}^+)^n} ds_1 \dots ds_n \mathcal{R}^n e^{-\mathcal{R}S(n)} \Pi_{m=1}^n |1 - \kappa(x_{S(m)})| r_{(x_{S(n)}, k_{S(n)})}(a), \quad (5.3)$$

and $S(m) = s_1 + \dots + s_m$.

Let $|k| > 4\bar{V}^{\frac{1}{2}}$ so that with two applications of Lemma 5.1, $\sup_{s, t \geq 0} |k_t - k_s| \leq |k|^{-1} 2\bar{V}$. In particular, for any time $S(n)$, we can apply Part (1) to the difference $|r_{(x_{S(n)}, k_{S(n)})}(a) - 1|$ to get

$$\begin{aligned} & \left| \tilde{r}_{(x, k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - \bar{\kappa} \sum_{n=0}^{\infty} \int_{(\mathbb{R}^+)^n} ds_1 \dots ds_n \mathcal{R}^n e^{-\mathcal{R}S(n)} \Pi_{m=1}^n |1 - \kappa(x_{S(m)})| \right| \\ & \leq \left(\frac{2\mathcal{R}}{|k|} + O\left(\frac{1}{|k|^2}\right) \right) \sum_{n=1}^{\infty} \int_{(\mathbb{R}^+)^n} ds_1 \dots ds_n \mathcal{R}^n e^{-\mathcal{R}S(n)} \Pi_{m=1}^n |1 - \kappa(x_{S(m)})| \\ & \leq \left(\frac{2\mathcal{R}}{|k|} + O\left(\frac{1}{|k|^2}\right) \right) \sum_{n=1}^{\infty} (1 - \nu)^n = \nu^{-1} (1 - \nu) \left(\frac{2\mathcal{R}}{|k|} + O\left(\frac{1}{|k|^2}\right) \right), \quad (5.4) \end{aligned}$$

where the second inequality follows since $|1 - \kappa(x_{S(m)})| \leq 1 - \nu$ for all m and $\mathcal{R}^n e^{-\mathcal{R}S(n)}$ defines a probability measure on $(\mathbb{R}^+)^n$.

If $\int_{(\mathbb{R}^+)^n} ds_1 \dots ds_n \mathcal{R}^n e^{-\mathcal{R}S(n)} \Pi_{m=1}^n |1 - \kappa(x_{S(m)})|$ were replaced by $(1 - \bar{\kappa})^n$ in the left-side of (5.4), then identity $\sum_{n=0}^{\infty} (1 - \bar{\kappa})^n = \kappa^{-1}$ would make the difference zero.

Using a telescoping sum and the definition of $r_{(x, k)}(a)$,

$$\begin{aligned} & \left| \int_{(\mathbb{R}^+)^n} ds_1 \dots ds_n \mathcal{R}^n e^{-\mathcal{R}S(n)} \Pi_{m=1}^n |1 - \kappa(x_{S(m)})| - (1 - \bar{\kappa})^n \right| \\ & \leq \sum_{m=0}^n (1 - \kappa)^{n-m} \int_{(\mathbb{R}^+)^{m-1}} ds_1 \dots ds_{m-1} \mathcal{R}^{m-1} e^{-\mathcal{R}S(m-1)} \Pi_{r=1}^{m-1} |1 - \kappa(x_{S(r)})| \\ & \quad \times \int_0^1 da |1 - \kappa(a)| |r_{(x_{S(m-1)}, k_{S(m-1)})}(a) - 1| \end{aligned}$$

Again by Part (1), since $\mathcal{R}^{m-1} e^{-\mathcal{R}S(m-1)}$ is a probability measure on $(\mathbb{R}^+)^{m-1}$, and by the bounds $1 - \kappa(a)$, $1 - \bar{\kappa} \leq 1 - \nu$, we can estimate the right-side above by

$$n(1 - \nu)^n \left(\frac{2\mathcal{R}}{|k|} + O\left(\frac{1}{|k|^2}\right) \right).$$

Putting everything together

$$\sup_{a \in [0, 1]} \left| \tilde{r}_{(x, k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \leq \left(\frac{2\mathcal{R}}{|k|} + O\left(\frac{1}{|k|^2}\right) \right) \sum_{n=0}^{\infty} (n+1)(1 - \nu)^n,$$

and the sum of the series is ν^{-2} .

Part (3):

Now we study the probability density $P_{(x,k)}(v)$ for the next momentum jump. We can write the density for the next momentum jump as

$$P_{(x,k)}(v) = \int_0^1 da \tilde{r}_{(x,k)}(a) \mathcal{P}_a(v), \quad (5.5)$$

We then have a bound using

$$\begin{aligned} \sup_{v \in \mathbb{R}} \left| \frac{P_{(x,k)}(v)}{\tilde{P}(v)} - 1 \right| &\leq \sup_{v \in \mathbb{R}} \frac{1}{\tilde{P}(v)} \int_0^1 da \left| \tilde{r}_{(x,k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \frac{\kappa(a)}{\bar{\kappa}} \mathcal{P}_a(v) \\ &\leq \sup_{a \in [0,1]} \left| \tilde{r}_{(x,k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \sup_{|v| \leq J} \frac{1}{\tilde{P}(v)} \int_0^1 da \mathcal{P}_a(v) \frac{\bar{\kappa}}{\kappa(a)} = \sup_{a \in [0,1]} \left| \tilde{r}_{(x,k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right|, \end{aligned} \quad (5.6)$$

where in the last equality we have used the definition of $\tilde{P}(v)$. Applying Part (2) then we get the bound.

Part (4): The density $T_{(x,k),t}$ can be written

$$T_{(x,k),t}(a) = n_{(x,k)}(a, t) \mathbf{k}_{(x,k)}(a)$$

where $\mathbf{k}_{(x,k)}(a)$ is defined as in Part (1), and $n_{(x,k)}(a, t)$ is the number of times the particle passes over the torus point a over a time period t when starting from (x, k) . The result follows by bounding the errors for $\mathbf{k}_{(x,k)}(a) \sim k$ and $|k|^{-1} n_{(x,k)}(a, t) \sim t$ which is similar to Part (1). $\square$

The following lemma bounds the contribution of the cumulative drift over periods of high-energy. Define $\tau(t)$ to be the time of the next to last momentum jump and put it equal to zero if two jumps have not occurred.

Lemma 5.3. *Assume List 2.1. In the limit $t \rightarrow \infty$, for $0 < \beta < \frac{1}{2}$,*

$$\mathbb{E} \left[\sup_{0 \leq r \leq t} \left| t^{-1+2\beta} \int_0^r ds \frac{dV}{dx}(X_s) \chi(|K_{\tau(s)}| > t^\beta) \right|^2 \right]^{\frac{1}{2}} \rightarrow 0.$$

Proof. It is convenient to split the total integral into a sum of integrals over the periods between Poisson times (at which there may be a momentum jump breaking the conservation of energy) which include only the even and odd terms respectively. Let $t_1, \dots, t_{N_r}$ be the Poisson times up to a time r .

$$\begin{aligned} \int_0^r ds \frac{dV}{dx}(X_s) \chi(|K_{\tau(s)}| > t^\beta) &\approx \sum_{n=1}^{\lfloor \frac{N_r}{2} \rfloor} \chi(|K_{t_{2n-1}}| \geq t^\beta) \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \\ &\quad + \sum_{n=0}^{\lfloor \frac{N_r}{2} \rfloor} \chi(|K_{t_{2n}}| \geq t^\beta) \int_{t_{2n+1}}^{t_{2n+2}} ds \frac{dV}{dx}(X_s), \end{aligned} \quad (5.7)$$

where on the right side, we have neglected the integral from $t_{2\lfloor \frac{N_r}{2} \rfloor}$ to r , which will be small, and we define $t_{-1} = 0$. We will focus on the sum with interval starting at even numbered times $[t_{2n}, t_{2n+1}]$. First, we will show that the terms in the sum can be replaced by the same

expressions conditioned on the information up to time t_{n-1} . These conditional expectations can be uniformly bounded by an argument following at the end of the proof. The strategy is to invoke Lemma 3.4 to guarantee that most of the terms in the sum have $|K_{t_{2n-1}}|$ on the order of $t^{\frac{1}{2}}$ rather than just $|K_{t_{2n}}| \geq t^\beta$. This lowers the bounds available for the conditional expectations of those terms.

The following is a martingale

$$Y_r = \sum_{n=0}^{\lfloor \frac{\mathcal{N}_t}{2} \rfloor} \chi(|K_{t_{2n-1}}| \geq \frac{1}{2}t^\beta) \left(\int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) - \mathbb{E} \left[\int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \middle| \mathcal{F}_{t_{2n-1}} \right] \right). \quad (5.8)$$

However, the second moment for a single term from the sum is bounded by

$$\begin{aligned} \mathbb{E} \left[\chi(|K_{t_{2n-1}}| \geq \frac{1}{2}t^\beta) \left| \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) - \mathbb{E} \left[\int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \middle| \mathcal{F}_{t_{2n-1}} \right] \right|^2 \right] \\ \leq \mathbb{E} \left[\left| \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \right|^2 \middle| |K_{t_{2n-1}}| \geq t^\beta \right] \leq 16\bar{V}^2 t^{-2\beta} \\ + 2\mathcal{R}^{-2} \sup_{0 \leq s \leq 1} \left| \frac{dV}{dx}(a) \right|^2 \Pr \left[|K_{t_{2n}}| \leq \frac{1}{2}t^{2\beta} \middle| |K_{t_{2n-1}}| \geq t^\beta \right], \quad (5.9) \end{aligned}$$

where we have considered separate bounds for the event that $|K_{t_{2n}}| \geq \frac{1}{2}t^\beta$ or $|K_{t_{2n}}| < \frac{1}{2}t^\beta$. When $|K_{t_{2n}}| \geq \frac{1}{2}t^\beta$, then we can apply Lemma 5.1 to bound the drift by $4\bar{V}t^{-\beta}$, and when $|K_{t_{2n}}| < \frac{1}{2}t^\beta$ then we use that the forces $|\frac{dV}{dx}(a)|$ are bounded and that the difference between two Poisson times has an exponential distribution with second moment $2\mathcal{R}^{-2}$. Finally, since the force can only change the momentum by at most $\bar{V}^{\frac{1}{2}} \ll \frac{1}{4}t^\beta$ over any time interval, only a large momentum jump can send $|K_{t_{2n}}|$ below $\frac{1}{2}t^\beta$. However, since the fourth moments of $\mathcal{P}_a(w)$ are less than ρ ,

$$\Pr \left[|K_{t_{2n}}| \leq \frac{1}{2}t^\beta \middle| |K_{t_{2n-1}}| \geq t^\beta \right] \leq \Pr \left[|w_n| \geq \frac{1}{4}t^\beta \middle| |K_{t_{2n-1}}| \geq t^\beta \right] \leq \sup_{a \in [0,1]} \int_{\frac{1}{4}t^\beta}^{\infty} dw \mathcal{P}_a(w) \leq \frac{4^4 \rho}{t^{4\beta}},$$

where the last inequality is Chebyshev's and thus the right side is $O(t^{-4\beta})$ which make the right term on the left side of (5.9) negligible compared to the left term.

Consider again the variance of a single term in (5.8). By Doob's maximal inequality

$$\mathbb{E} \left[\sup_{0 \leq r \leq t} |Y_r|^2 \right] \leq 4 \mathbb{E} [|Y_t|^2] \leq (16\bar{V}^2 t^{-2\beta} + O(t^{-4\beta})) \mathbb{E} [\mathcal{N}_t] = 16\bar{V}^2 \mathcal{R} t^{1-2\beta} + O(t^{1-4\beta}).$$

Thus we can focus on bounding the expressions $|\mathbb{E} [\int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \middle| \mathcal{F}_{t_{2n-1}}]|$ when $|K_{t_{2n-1}}| \geq t^\beta$. The end result of the analysis below will be to show that there is a constant c such that for all sufficiently large $t^\beta \gg 1$

$$\left| \mathbb{E} [\chi(|K_{t_{2n-1}}| \geq t^\beta) \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \middle| \mathcal{F}_{t_{2n-1}}] \right| \leq c t^{-2\beta}. \quad (5.10)$$

Applying the above inequality (also with t^β replaced by $\epsilon t^{\frac{1}{2}}$),

$$\sum_{n=1}^{\lfloor \frac{\mathcal{N}_t}{2} \rfloor} \left| \mathbb{E} [\chi(|K_{t_{2n-1}}| \geq t^\beta) \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \middle| \mathcal{F}_{t_{2n-1}}] \right| \leq c \mathcal{N}_{t,\epsilon} \epsilon^{-2} t^{-1} + c (\mathcal{N}_t - \mathcal{N}_{t,\epsilon}) t^{-2\beta},$$

where $\mathcal{N}_{r,\epsilon}$ is the number of terms with $|K_{t_{2n-1}}| \geq \epsilon t^{\frac{1}{2}}$ up to time r . By the triangle inequality

$$\begin{aligned} & \mathbb{E} \left[\left| t^{-1+2\beta} \sum_{n=1}^{\lfloor \frac{\mathcal{N}_t}{2} \rfloor} \mathbb{E} [\chi(|K_{t_{2n-1}}| \geq t^\beta) \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \mid \mathcal{F}_{t_{2n-1}}] \right|^2 \right]^{\frac{1}{2}} \\ & \leq 2^{\frac{1}{2}} c \mathcal{R} \epsilon^{-2} t^{-1+2\beta} + c t^{-1} \mathbb{E} [|\mathcal{N}_t - \mathcal{N}_{t,\epsilon} - \gamma_\epsilon(t)|^2]^{\frac{1}{2}} + c t^{-1} \mathbb{E} [|\gamma_\epsilon(t)|^2]^{\frac{1}{2}}, \quad (5.11) \end{aligned}$$

where it was used that $\mathcal{N}_{t,\epsilon} \leq \mathcal{N}_t$, $\mathbb{E} [|\mathcal{N}_t|^2]^{\frac{1}{2}} \leq 2^{\frac{1}{2}} t \mathcal{R}$, and $\gamma_\epsilon(t)$ is defined as

$$\gamma_\epsilon(t) = \mathcal{R} \sum_{n=1}^{\mathcal{N}_r} (t_n - t_{n-1}) \chi(|K_{t_{n-1}}| \leq \epsilon t^{\frac{1}{2}}).$$

The difference $\mathcal{N}_r - \mathcal{N}_{t,\epsilon} - \gamma_\epsilon(r)$ is a martingale since $t_n - t_{n-1}$ are exponentially distributed with mean $\mathcal{R}^{-1}$. The variance of the martingale satisfies

$$\mathbb{E} [|\mathcal{N}_t - \mathcal{N}_{t,\epsilon} - \gamma_\epsilon(t)|^2] = \mathbb{E} \left[\sum_{n=1}^{\mathcal{N}_t} |\mathcal{R}(t_n - t_{n-1}) - 1|^2 \chi(|K_{t_{n-1}}| \leq \epsilon t^{\frac{1}{2}}) \right] \leq \mathbb{E} [\mathcal{N}_t].$$

Thus the middle term on the right-side of (5.11) is $O(t^{\frac{1}{2}-2\beta})$. $\gamma_\epsilon(t)$ is less than the amount of time $r \in [0, t]$ the particle spends with $t^{-\frac{1}{2}} E_r^{\frac{1}{2}} \leq \epsilon$. In other terms, $t^{-1} \gamma_\epsilon(t) \leq 1 - T_{\epsilon,V}^{(t)}$ where $T_{\epsilon,V}^{(t)}$ is defined as in Lemma 3.4. Thus by Lemma 3.4, $\Pr [t^{-1} \gamma_\epsilon(t) \geq \delta] \leq C \frac{r_2^{\frac{1}{2}}}{r_1} \frac{\epsilon}{\delta}$. Since $t^{-1} \gamma_\epsilon(t)$ is bounded by 1,

$$\mathbb{E} [|t^{-1} \gamma_\epsilon(t)|^2]^{\frac{1}{2}} \leq \delta \Pr [t^{-1} \gamma_\epsilon(t) < \delta] + \Pr [t^{-1} \gamma_\epsilon(t) \geq \delta] \leq \delta + C \frac{r_2^{\frac{1}{2}}}{r_1} \frac{\epsilon}{\delta}$$

Thus we can pick δ to make the first term small and then pick ϵ to make the second term small. We now turn to showing (5.10).

By the Markov property

$$\begin{aligned} & \left| \mathbb{E} [\chi(|K_{t_{2n-1}}| \geq t^\beta) \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \mid \mathcal{F}_{t_{2n-1}}] \right| \\ & \leq \int dx dk P_\omega(x, k) \chi(|k| \geq t^\beta) \left| \mathbb{E} \left[\int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \mid (X_{t_{2n-1}}, K_{t_{2n-1}}) = (x, k) \right] \right|. \quad (5.12) \end{aligned}$$

where $P_\omega(x, k)$ is the distribution $(X_{t_{2n-1}}, K_{t_{2n-1}})$ conditioned on $\omega \in \mathcal{F}_{t_{2n-1}}$.

Assuming that $|K_{t_{2n-1}}| \geq t^\beta$ and $K_{t_{2n}} \geq \frac{1}{2} t^\beta$,

$$\begin{aligned} & \left| \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) - \frac{V(X_{t_{2n+1}}) - V(X_{t_{2n}})}{K_{t_{2n-1}}} \right| \\ & \leq \left| \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) - \frac{1}{K_{t_{2n}}} \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) K_s \right| + \left| \frac{1}{K_{t_{2n}}} - \frac{1}{K_{t_{2n-1}}} \right| |V(X_{t_{2n+1}}) - V(X_{t_{2n}})| \\ & < 2t^{-2\beta} \bar{V} \sup_{0 \leq x \leq 1} \left| \frac{dV}{dx}(x) \right| (t_{2n+1} - t_{2n}) + 2t^{-2\beta} \bar{V} |K_{t_{2n}} - K_{t_{2n-1}}|, \quad (5.13) \end{aligned}$$

where we have used the identity $\int_r^t ds \frac{dV}{dx}(X_s) K_s = V(X_t) - V(X_r)$. The only thing random in the final bound is the difference $t_{n+1} - t_n$, which is an exponential random variable with mean $\mathcal{R}^{-1}$ and the difference $|K_{t_{2n}} - K_{t_{2n-1}}|$ which has variance less than $\frac{r_2^2}{\mathcal{R}}$. Thus

$$\begin{aligned} & \left| \mathbb{E} \left[\chi(|K_{t_{2n-1}}| \geq t^\beta) \int_{t_{2n}}^{t_{2n+1}} ds \frac{dV}{dx}(X_s) \middle| \mathcal{F}_{t_{2n-1}} \right] \right| \\ & \leq O(t^{-4\beta}) + 2t^{-2\beta} \mathcal{R} \bar{V} \sup_{0 \leq x \leq 1} \left| \frac{dV}{dx}(x) \right| + 2t^{-2\beta} \bar{V} \frac{r_2^{\frac{1}{2}}}{\mathcal{R}^{\frac{1}{2}}} \\ & + \int dx dk P_\omega(x, k) \chi(|k| \geq t^\beta) \left| \mathbb{E} \left[\frac{V(X_{t_{2n+1}}) - V(X_{t_{2n}})}{K_{t_{2n-1}}} \middle| (X_{t_{2n-1}}, K_{t_{2n-1}}) = (x, k) \right] \right|, \quad (5.14) \end{aligned}$$

where $\omega \in \mathcal{F}_{t_{2n-1}}$ and $O(t^{-4\beta})$ corresponds the unlikely event that $\Pr[K_{t_{2n}} < \frac{1}{2}t^\beta]$ which we have treated above following (5.9).

Adding and subtracting the spatial average of the potential, $\int_0^1 da V(a) = \mathcal{V}$ in the expectation above,

$$\begin{aligned} & \left| \mathbb{E} \left[\frac{V(X_{t_{2n+1}}) - V(X_{t_{2n}})}{K_{t_{2n-1}}} \middle| (X_{t_{2n-1}}, K_{t_{2n-1}}) \right] \right| \\ & \leq \frac{1}{|K_{t_{2n-1}}|} \left(\int_{\mathbb{R}^2} dx dk P_{(X_{t_{2n-1}}, K_{t_{2n-1}})}(x, k) \left| \mathbb{E}[V(X_{t_{2n+1}}) - \mathcal{V} \mid (X_{t_{2n}}, K_{t_{2n}}) = (x, k)] \right| \right. \\ & \quad \left. + \left| \mathbb{E}[V(X_{t_{2n}}) - \mathcal{V} \mid (X_{t_{2n-1}}, K_{t_{2n-1}})] \right| \right), \quad (5.15) \end{aligned}$$

where $P_{(X_{t_{2n-1}}, K_{t_{2n-1}})}$ is the probability density for $(X_{t_{2n}}, K_{t_{2n}})$ given $(X_{t_{2n-1}}, K_{t_{2n-1}})$.

Finally, we can work with quantities that allow more explicit expressions

$$\mathbb{E}[V(X_{t_{2n}}) \mid (X_{t_{n-1}}, K_{t_{n-1}})] = \int_0^\infty dt \mathcal{R} e^{-\mathcal{R}t} V(x_t) = \int_0^1 da r_{(x_0, k_0)}(a) V(a),$$

where x_t is the position at time t for the particle evolving according to the dynamics from the initial point $(x_0, k_0) = (X_{t_{n-1}}, K_{t_{n-1}})$, and $r_{(x, k)} \in L^1([0, 1])$ is defined as in Lemma 5.2.

$$\left| \mathbb{E}[V(X_{t_{2n}}) \mid (X_{t_{n-1}}, K_{t_{n-1}})] - \mathcal{V} \right| \leq \bar{V} \int_0^1 da |r_{(x_0, k_0)}(a) - 1|.$$

By Part (1) of Lemma 5.2, when $|K_{t_{n-1}}| \geq t^\beta$, then $|r_{(x_0, k_0)}(a) - 1| \leq 4\mathcal{R}t^{-\beta} + O(t^{-2\beta})$. A similar analysis bounds the term $\left| \mathbb{E}[V(X_{t_{2n+1}}) - \mathcal{V} \mid (X_{t_{2n}}, K_{t_{2n}}) = (x, k)] \right|$. Thus with the factor of $|K_{t_{n-1}}|^{-1}$ on the right side of (5.15), then (5.15) is $O(t^{-2\beta})$, which completes the proof. $\square$

6 Bounding the momentum drift

In general, we have that $K_t = K_0 + M_t + \int_0^t dr \frac{dV}{dx}(X_r)$. In this section, we develop tools for controlling the cumulative drift $\int_0^{st} dr \frac{dV}{dx}(X_r)$. The end result, under the assumption of the symmetry (iii) of 2.2, is that

$$\mathbb{E} \left[\left| \sup_{0 \leq s \leq 1} t^{-\frac{1}{2}} \int_0^{st} ds \frac{dV}{dx}(X_s) \right| \right] \longrightarrow 0. \quad (6.1)$$

Thus on the scale $t^{\frac{1}{2}}$ of a central limit theorem for K_t , the drift term vanishes.

By the Lemma 3.4, the particle spends most of the time at “high energy” (in Lemma 3.4 this meant $\propto t$), where the contribution to the total drift over any given finite time interval is small. However, the particle is also making occasional shorter incursions to “low energy” where the contribution may be larger over a finite interval. In this section, “low energy” roughly means below $t^{\frac{1}{4}}$. In order to bound (6.1), the analysis is split into parts treating the drift at high and low energies respectively.

From this section onwards, we contract the position degree of freedom to a single periodic cell $x \in [0, 1]$. This clearly does not affect the statistics for the drift process (6.1). Thus the dynamics satisfies the same linear Boltzmann equation (2.1) as before but with periodic boundary conditions; the derivatives in position at the boundaries of the interval are symmetric.

We now define what we mean by low energy incursions. They are limited by starting and ending times. Define the hitting time $\theta_0 = \min\{s \in [0, t] \mid |K_s| \geq t^{\frac{1}{4}}\}$. For $j \geq 1$ define the sequences of hitting times σ_j, θ_j :

$$\sigma_j = \min\{s \in [0, \infty), M_s - M_{s-} \neq 0 \mid s > \theta_{j-1}, |K_s| < t^{\frac{1}{4}}\}, \quad (6.2)$$

$$\theta_j = \min\{s \in [0, \infty) \mid s > \sigma_j, |K_s| > 2t^{\frac{1}{4}}\}, \quad (6.3)$$

Notice that θ_0 is defined differently than θ_j for $j \geq 1$. We refer to $[\sigma_j, \theta_j]$ as the time period of the j th incursion.

In the lemma below, we give a bound on the expected number of incursions $N_Y(\varrho)$ over a time interval $[0, \varrho t]$, and show that the time periods of incursions $\theta_j - \sigma_j$ have finite first moments. The time periods between incursions $\sigma_{j+1} - \theta_j$ can be shown to be almost surely finite. This follows from an argument using Lévy’s zero-one law and Theorem 2.3, but showing $\sigma_{j+1} - \theta_j$ to be finite is not required to prove Theorem 2.3. In any case, bounds on $\sigma_{j+1} - \theta_j$ are intrinsically less important to us, since the challenge is to get estimates for the low-energy part of the walk.

Lemma 6.1. *Assume 2.1 and (i) of List 2.2.*

1. *Given $\sigma_j < \infty$, the difference $\theta_j - \sigma_j$ has expectation $O(t^{\frac{1}{2}})$.*
2. *Let $\varrho > 0$. For large enough t , the expectation for the number of incursions in the interval $[0, \varrho t]$ is bounded as*

$$\mathbb{E}[N_Y(\varrho t)] \leq \varrho^{\frac{1}{2}} r_2^{\frac{1}{2}} t^{\frac{1}{4}}$$

Proof.

Part (1):

Let us set construct the stopping time $\theta_{\mathcal{T}} = (\theta_j - \sigma_j) \wedge \mathcal{T}$ for some bound $\mathcal{T} > 1$ and set $\sigma_j = 0$.

$$\mathbb{E}[\theta_{\mathcal{T}}] \leq \frac{r_2}{r_1} \mathbb{E}[\langle M \rangle_{\theta_{\mathcal{T}}}] = \frac{r_2}{r_1} \mathbb{E}[[M]_{\theta_{\mathcal{T}}}],$$

where the equality follows since $[M]_t - \langle M \rangle_t$ is a martingale, and the inequality is a consequence of $r_1 \leq \frac{d}{dt} \langle M \rangle_t \leq r_2$.

Lemma 3.5 states that $2^{-1} \sum_{n=1}^{\mathcal{N}_r} w_n^2 = 2^{-1} [M]_r$ differs from $E_r - E_0$ by a martingale, and by the Optional Sampling Theorem the expectation of the martingale part is zero at time $\theta_{\mathcal{T}}$ so

$$\mathbb{E}[[M]_{\theta_{\mathcal{T}}}] = 2\mathbb{E}[E_{\theta_{\mathcal{T}}} - E_0].$$

Let D be the size of the over-jump of the boundary $-2t^{\frac{1}{4}}$ or $2t^{\frac{1}{4}}$ when $\theta_{\mathcal{T}} < \mathcal{T}$, and $\bar{V}$ be the max of the potential.

$$\mathbb{E}[E_{\theta_{\mathcal{T}}} - E_0] \leq \mathbb{E}\left[\frac{1}{2}(2t^{\frac{1}{4}} + D)^2 + \bar{V}|\theta_{\mathcal{T}} < \mathcal{T}\right] + (2t^{\frac{1}{2}} + \bar{V}) \Pr[\theta_{\mathcal{T}} = \mathcal{T}] \leq 2t^{\frac{1}{2}} + O(1) < 3^{\frac{1}{2}}$$

By Lemma B.4, there are universal bounds determined by $\mathcal{C}$ and η on all the moments of D , $\mathbb{E}[D^2] < \rho_2(\mathcal{C}, \eta)$. Using that $|x + y|^2 \leq 2x^2 + 2y^2$, $\Pr[\theta > \mathcal{T}] \leq 1$, and that $\bar{V} \ll t^{\frac{1}{4}}$

$$\mathbb{E}[\theta_{\mathcal{T}}] < \frac{r_2}{r_1}(6t^{\frac{1}{2}} + O(1)) = O(t^{\frac{1}{2}}).$$

Finally, by taking the limit $\mathcal{T} \rightarrow \infty$, we get a bound for the second moment of $\theta_j - \sigma_j$:

$$\mathbb{E}[\theta_j - \sigma_j] = \limsup_{\mathcal{T} \rightarrow \infty} \mathbb{E}[\theta_{\mathcal{T}}] \leq \frac{r_2}{r_1}(6t^{\frac{1}{2}} + O(1)) = O(t^{\frac{1}{2}}).$$

Part (2):

By Part (1) each incursion ends. Thus for each count of $N_Y(\varrho t)$ there is a distinct up-crossing in which $|K_s|$ begins below $t^{\frac{1}{4}}$ and ends up above $2t^{\frac{1}{4}}$. However, for large values of momentum $\frac{1}{\sqrt{2}}|K_s| \approx E_s^{\frac{1}{2}}$, and we bound N_Y by the number of up-crossings $U_{\varrho t}(2^{-1}t^{\frac{1}{4}}, 2t^{\frac{1}{4}}; E_s(\omega))$ that $E_s^{\frac{1}{2}}$ makes between $2^{-1}t^{\frac{1}{4}}$ and $2t^{\frac{1}{4}}$. Since $E_s^{\frac{1}{2}}$ is a submartingale, we can apply the submartingale up-crossing inequality [3] to attain

$$\mathbb{E}[N_Y(\varrho t)] \leq \mathbb{E}[U_{\varrho t}(2^{-1}t^{\frac{1}{4}}, 2t^{\frac{1}{4}}; E_s(\omega))] \leq \frac{\mathbb{E}[E_{\varrho t}^{\frac{1}{2}}]}{2t^{\frac{1}{4}} - 2^{-1}t^{\frac{1}{4}}} \approx \frac{\sqrt{2}}{3} r_2^{\frac{1}{2}} \varrho^{\frac{1}{2}} t^{\frac{1}{4}} < r_2^{\frac{1}{2}} \varrho^{\frac{1}{2}} t^{\frac{1}{4}}.$$

□

The basic idea for our analysis is the following:

- For large t , there is an asymptotic independence between the events during a single incursion and all events up to the end of a previous incursion;
- Events during the incursion, which occur far enough after the starting time of the incursion are independent of the initial state of the incursion.

It is convenient in many places to have an effective bound on the size of the momentum jumps that are likely to occur in the interval $[0, t]$. We thus consider the statistics for our model conditioned on the event

$$\{|v_n| \leq t^{\frac{1}{40}} \text{ for all } n \text{ such that } t_n \leq t\},$$

where a jump greater than $t^{\frac{1}{40}}$ is considered to be large. The lemma below shows that the probability that there is a jump above $t^{\frac{1}{40}}$ over the interval $[0, t]$ decays super-polynomially, and that for dealing with the drift over up to time t , we can neglect the possibility of large jumps. The choice of $\frac{1}{40}$ involves constraints from Proposition B.6.

Lemma 6.2. Assume (i) of List 2.2, then the probability of a momentum jump v_n with $|v_n| \geq t^{\frac{1}{40}}$ over the interval $[0, t]$ is $O(t e^{-\frac{\eta}{2} t^{\frac{1}{40}}})$, for η as in (i) of List 2.2. Moreover, the difference in the quantity

$$\mathbb{E} \left[\left| \sup_{0 \leq s \leq 1} t^{-\frac{1}{2}} \int_0^{st} ds \frac{dV}{dx}(X_s) \right|^2 \right]^{\frac{1}{2}},$$

for the dynamics conditioned not to make jumps greater than $t^{\frac{1}{40}}$ and the unconditioned dynamics is $O(t^2 e^{-\frac{\eta}{2} t^{\frac{1}{40}}})$

Proof. Assumption (i) of List 2.2 implies that the Laplace transforms for single momentum jumps are uniformly bounded by $\mathcal{C}(1 - e^{-\eta-q})^{-1}$. Using Chebyshev's inequality we can bound the probability that any jump is greater than $t^{\frac{1}{40}}$. On the other hand, the integral of the drift can be at most $t \sup_{a \in [0,1]} \left| \frac{dV}{dx}(a) \right|$. □

We refer to Appendix B for a discussion of boundary crossing distributions and the definition of the boundary crossing density $\phi_\infty : [0, 1] \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$. If H is a random variable, then $\Pr[H = y]$, for dummy variable $y \in \mathbb{R}$, refers to the distributional measure of H or its probability density if it exists. For a signed measure μ on $\mathbb{R}$, then $\|\mu\|_1 = |\mu|(\mathbb{R})$, where $|\mu|$ is the absolute value of the measure. Most of the random variables in this article (e.g. energy and momentum jumps) have well-defined densities.

The following lemma states that incursions beginning at points $(X_{\sigma_j}, K_{\sigma_j})$ for $s_1 K_{\sigma_j} \in [t^{\frac{1}{4}} - t^{\frac{1}{40}}, t^{\frac{1}{4}}]$ for fixed $s_1 = \pm$ all have approximately the same probabilities for ending the incursion in the positive or negative direction.

Lemma 6.3. Assume List 2.1 and (i)-(ii) of List 2.2. Consider the dynamics conditioned not to have jumps greater than $t^{\frac{1}{40}}$. Let $\mathbf{s}_1, \mathbf{s}_2 \in \{+, -\}$, and $(x, \mathbf{s}_1 k) \in [0, 1] \times [t^{\frac{1}{4}} - t^{\frac{1}{40}}, t^{\frac{1}{4}}]$. There are constants $\rho_{\mathbf{s}_1, \mathbf{s}_2}(t)$ such that as $t \rightarrow \infty$

$$\sup_{(x,k)} \left| \frac{\Pr[\mathbf{s}_2 K_{\theta_j} > 0 \mid (X_{\sigma_j}, \mathbf{s}_1 K_{\sigma_j}) = (x, k)]}{\rho_{\mathbf{s}_1, \mathbf{s}_2}(t)} - 1 \right| \rightarrow 0$$

Proof. Fix $\mathbf{s}_1 = \mathbf{s}_2 = +$. Define $\phi_{\uparrow, t}^{(a, v)}, \phi_{\downarrow, t}^{(a, v)}$ to be the boundary crossing distributions above and below the set $S = [t^{\frac{1}{4}} - t^{\frac{1}{40}}, t^{\frac{1}{4}} + t^{\frac{1}{40}}]$ starting from $(a, t^{\frac{1}{4}} - v)$.

Using the Markov property and that H is a function of the process after the time τ ,

$$\begin{aligned} & \Pr[K_{\theta_j} > 0 \mid (X_{\sigma_j}, K_{\sigma_j}) = (a, t^{\frac{1}{4}} - v)] \\ &= \int_{[0,1] \times \mathbb{R}^+} dq dp \left(\phi_{\uparrow, t}^{(a, v)}(q, p) \Pr[K_{\theta_j} > 0 \mid (X_\tau, K_\tau) = (q, t^{\frac{1}{4}} + t^{\frac{1}{20}} + p)] \right. \\ & \quad \left. + \phi_{\downarrow, t}^{(a, v)}(q, p) \Pr[K_{\theta_j} > 0 \mid (X_\tau, K_\tau) = (q, t^{\frac{1}{4}} - t^{\frac{1}{20}} - p)] \right)_1. \end{aligned} \quad (6.4)$$

Thus for $(a, v), (a', v') \in [0, 1] \times [0, t^{\frac{1}{40}}]$,

$$\begin{aligned} & \sup_{a, a', v, v'} \left\| \Pr[H = y \mid (X_{\sigma_j}, K_{\sigma_j}) = (a, t^{\frac{1}{4}} - v)] - \Pr[H = y \mid (X_{\sigma_j}, K_{\sigma_j}) = (a', t^{\frac{1}{4}} - v')] \right\|_1 \\ & \leq \sup_{a, a', v, v'} \left\| \phi_{\uparrow, t}^{(a, v)} - \phi_{\uparrow, t}^{(a', v')} \right\|_1 + \sup_{a, a', v, v'} \left\| \phi_{\downarrow, t}^{(a, v)} - \phi_{\downarrow, t}^{(a', v')} \right\|_1 \end{aligned} \quad (6.5)$$

By Proposition B.7, $\phi_{\uparrow,t}^{(a,v)}$ and $\phi_{\downarrow,t}^{(a,v)}$ converge uniformly to ϕ_∞ in L^1 . Thus the diameter $D(t)$ of the set of possible values for $\Pr [K_{\theta_j} > 0 \mid (X_\tau, K_\tau) = (a, t^{\frac{1}{4}} - v)]$ as a function of (a, v) shrinks to zero as $t \rightarrow \infty$. Let us define

$$\rho_{+,+}(t) = \Pr [K_{\theta_j} > 0 \mid (X_{\sigma_j}, K_{\sigma_j}) = (x, k)] + D_t^{\frac{1}{2}} + \frac{1}{t},$$

for any choice of (x, k) where t^{-1} is merely to ensure that $\rho_{+,+}(t)$ is non-zero, and the square root is introduced so that $D_t D_t^{-\frac{1}{2}} = D_t^{\frac{1}{2}} \rightarrow 0$. Then we will have the conclusion of the lemma. $\square$

6.1 Bounding the drift over an incursion

In this section, we define incursions to have end times ς_j which are different but related to the end times to θ_j . Define the sequence of times ς_j :

$$\varsigma_j = \min\{s \in [0, t], M_s - M_{s-} \neq 0 \mid s > \sigma_j, \inf_{s < r \leq \theta_j} |K_r| > t^{\frac{1}{4}}\}. \quad (6.6)$$

The ς_j 's are not hitting times since information up to time θ_j is required to determine them. However, for the time-reversed dynamics the ς_j are well-defined hitting times and are defined to be symmetric with the σ_j 's. This will be important in the next section. All of the results in this section apply when ς_j is replaced by θ_j .

Define the random variables

$$Y_j = t^{-\frac{1}{4}} \int_{\sigma_j}^{\varsigma_j} dr \frac{dV}{dx}(X_r).$$

We consider the Y_r 's to contain the low energy contribution to the cumulative drift. Yet, during some portion of the interval $[\sigma_j, \varsigma_j]$, the particle is likely to spend some time with momentum above $t^{\frac{1}{4}}$, and thus have overlap with the quantity in Lemma 5.3 with $\beta = \frac{1}{4}$. Let $N_Y(t)$ be the number of Y_j terms up to time t . It is convenient to split the Y_j 's into four classes. For $m \geq 1$, $\mathbf{s}_1, \mathbf{s}_2 \in \{+, -\}$,

$$Y_{\mathbf{s}_1, \mathbf{s}_2}(m) = t^{-\frac{1}{4}} \chi(\mathbf{s}_1 K_{\sigma_j} > 0, \mathbf{s}_2 K_{\varsigma_j} > 0) \int_{\sigma_j}^{\varsigma_j} dr \frac{dV}{dx}(X_r),$$

where j and m are related through

$$j = \min\{n \geq 0 \mid m = \sum_{i=1}^n \chi(\mathbf{s}_1 K_{\sigma_i} > 0)\}. \quad (6.7)$$

$Y_{\mathbf{s}_1, \mathbf{s}_2}(m)$ is equal to the drift for the m th incursion that begins with momentum having sign s_1 provided that the incursion ends with sign s_2 . Naturally, if there is no m th incursion with sign s_1 , then we set $Y_{\mathbf{s}_1, \mathbf{s}_2}(m) = 0$. For $\mathbf{s} \in \{\pm\}$, we also define $N_{\mathbf{s}}(r)$ to be the number $N_{\mathbf{s}}(r) = \sum_{i=1}^{N_Y(r)} \chi(\mathbf{s} K_{\sigma_i} > 0)$.

Define the constant

$$c_{\mathbf{s}_1, \mathbf{s}_2}(t) = t^{-\frac{1}{4}} \int_{[0,1] \times (0, t^{\frac{1}{40}})} da dv \phi(a, v) \mathbb{E}_{(a, \mathbf{s}_1 t^{\frac{1}{4}} - \mathbf{s}_1 v)} \left[\int_0^\varsigma ds \frac{dV}{dx}(X_s) \chi(\mathbf{s}_2 K_\varsigma > 0) \right], \quad (6.8)$$

where ς is defined analogously to the ς_j 's as the last time that there is a momentum jump inside $(-t^{\frac{1}{4}}, t^{\frac{1}{4}})$ before exiting the larger interval $(-2t^{\frac{1}{4}}, 2t^{\frac{1}{4}})$:

$$\begin{aligned}\varsigma &= \min\{s \in [0, \theta), M_s - M_{s-} \neq 0 \mid \inf_{s < r \leq \theta} |K_r| > t^{\frac{1}{4}}\}, \\ \theta &= \min\{s \in [0, \infty) \mid |K_s| > 2t^{\frac{1}{4}}\}.\end{aligned}$$

In fact, $c_{\mathbf{s}_1, \mathbf{s}_2}(t)$ will be asymptotically close to $t^{-\frac{1}{4}} \mathbb{E}_{(a, \mathbf{s}_1 t^{\frac{1}{4}} - \mathbf{s}_1 v)} \left[\int_0^\varsigma ds \frac{dV}{dx}(X_s) \chi(\mathbf{s}_2 K_\theta > 0) \right]$ for any fixed $(a, v) \in [0, 1] \times \mathbb{R}^+$ as $t \rightarrow \infty$ so the density $\phi(a, v)$ appearing in the definition of $c_{\mathbf{s}_1, \mathbf{s}_2}$ is not important (except as a matter of convenience).

The main purpose of the following proposition is to establish Part (3) which says that the sum of the $Y_{\mathbf{s}_1, \mathbf{s}_2}(m)$'s can be replaced by the number $N_{\mathbf{s}_1}(st)$ multiplied by the constant $c_{\mathbf{s}_1, \mathbf{s}_2}(t)$.

Proposition 6.4. *Assume List 2.1 and (i)-(ii) of 2.2.*

1. *For large enough t , then for all j and $\omega \in \mathcal{F}_{\sigma_j}$*

$$\mathbb{E}[Y_j^2 | \mathcal{F}_{\sigma_j}]^{\frac{1}{2}} < 5.$$

2. *Let $c_{\mathbf{s}_1, \mathbf{s}_2}(t)$ be defined as in (6.8) and j and m be related by (6.7). As $t \rightarrow \infty$, we have the $L^2(\Omega)$ convergence*

$$\mathbb{E} \left[\left| \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t) \right|^2 \right]^{\frac{1}{2}} \rightarrow 0.$$

3. *As $t \rightarrow \infty$, there is $L^2(\Omega)$ convergence*

$$t^{-\frac{1}{4}} \mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| \sum_{m=1}^{N_{\mathbf{s}_1}(st)} Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - N_{\mathbf{s}_1}(st) c_{\mathbf{s}_1, \mathbf{s}_2}(t) \right| \right] \rightarrow 0.$$

Proof.

Part (1):

By Lemma 6.2, we can take the jumps to be bounded by $t^{\frac{1}{40}}$, although we will not employ this till the end of the proof. Set $\sigma = \sigma_j$, $\varsigma = \varsigma_j$, and $\theta = \theta_j$. By the triangle inequality,

$$\mathbb{E}[Y_j^2 | \mathcal{F}_\sigma]^{\frac{1}{2}} \leq t^{-\frac{1}{4}} \mathbb{E}[(M_{\varsigma-} - M_\sigma)^2 | \mathcal{F}_\sigma]^{\frac{1}{2}} + t^{-\frac{1}{4}} \mathbb{E}[(K_{\varsigma-} - K_\sigma)^2 | \mathcal{F}_\sigma]^{\frac{1}{2}}.$$

The times σ and ς are defined such that $|K_\sigma|, |K_{\varsigma-}| \leq t^{\frac{1}{4}}$. Thus $\mathbb{E}[(K_{\varsigma-} - K_\sigma)^2 | \mathcal{F}_\sigma]^{\frac{1}{2}} \leq 2t^{\frac{1}{4}}$.

As in the proof of Part (1) of Lemma 6.1 define the stopping time $\theta_\mathcal{T} = \theta \wedge \mathcal{T}$ and the capped time $\varsigma_\mathcal{T} = \varsigma \wedge \mathcal{T}$.

By Doob's maximal inequality and $\varsigma_\mathcal{T} \leq \theta_\mathcal{T}$,

$$\begin{aligned}\mathbb{E}[|M_{\varsigma_\mathcal{T}-} - M_\sigma|^2 | \mathcal{F}_\sigma]^{\frac{1}{2}} &\leq \mathbb{E} \left[\sup_{\sigma \leq r \leq \theta_\mathcal{T}} |M_r - M_\sigma|^2 | \mathcal{F}_\sigma \right]^{\frac{1}{2}} \\ &\leq 2\mathbb{E}[|M_{\theta_\mathcal{T}} - M_\sigma|^2 | \mathcal{F}_\sigma]^{\frac{1}{2}} = 2\mathbb{E}[|M|_{\theta_\mathcal{T}} - |M|_\sigma | \mathcal{F}_\sigma]^{\frac{1}{2}}, \quad (6.9)\end{aligned}$$

where $[M]_t$ is the quadratic variation and of the martingale M_t up to time t , and the last equality follows from the optional sampling theorem. By Lemma 3.5, $\mathcal{A}_t$ in the increasing part in a Doob-Meyer decomposition for E_t ,

$$2\mathcal{A}_t - 2\mathcal{A}_s = [M]_t - [M]_s.$$

Applying the above for the time interval from σ up to stopping time θ_T , the right-hand side of (6.9), is bounded by

$$2\mathbb{E}[\mathcal{A}_{\theta_T} - \mathcal{A}_\sigma | \mathcal{F}_\sigma] = 2\mathbb{E}[E_{\theta_T} - E_\sigma | \mathcal{F}_\sigma] \leq 4t^{\frac{1}{2}} + O(1)$$

where the equality is another use of the optional sampling theorem. The inequality comes from the proof of Part (1) of Lemma 6.1.

Part (2):

Let us take $\mathbf{s}_1, \mathbf{s}_2 = +$. By Lemma 6.2, we can take jumps to be bounded by $t^{\frac{1}{40}}$. By the Markov property and by the definition of $Y_{+,+}(m)$,

$$\begin{aligned} \mathbb{E}[Y_{+,+}(m) | \mathcal{F}_{\theta_{j-1}}] &= \mathbb{E}_{(X_{\theta_{j-1}}, K_{\theta_{j-1}})} \left[t^{-\frac{1}{4}} \int_0^\varsigma ds \frac{dV}{dx}(X_s) \chi(K_\varsigma > 0) \right] \\ &= \int_{[0,1] \times [0, t^{\frac{1}{40}}]} da' dv' \phi_t(a', v') \mathbb{E}_{(a', t^{\frac{1}{4}} - v')} \left[t^{-\frac{1}{4}} \int_0^\varsigma ds \frac{dV}{dx}(X_s) \chi(K_\varsigma > 0) \right]. \end{aligned} \quad (6.10)$$

where $\phi_t(a, v)$ is the joint distribution of $(X_{\sigma_j}, -K_{\sigma_j} + t^{\frac{1}{4}})$ given $(X_{\theta_{j-1}}, K_{\theta_{j-1}})$. We attach a subscript (a, v) to the symbol ϕ_t to indicate the ending point $(X_{\theta_{j-1}}, K_{\theta_{j-1}}) = (a, t^{\frac{1}{4}} - v)$ of the last previous incursion.

We have that

$$\begin{aligned} &\mathbb{E} \left[\left| c_{+,+}(m) - \mathbb{E}[Y_{+,+}(m) | \mathcal{F}_{\theta_{j-1}}] \right|^2 \right] \\ &\leq \sup_{(a', v')} \mathbb{E}_{(a', t^{\frac{1}{4}} - v')} \left[\left| t^{-\frac{1}{4}} \int_0^\varsigma ds \frac{dV}{dx}(X_s) \right|^2 \chi(K_\varsigma > 0) \right] \mathbb{E} \left[\left\| \phi_{t, (a, v)} - \phi_\infty \right\|_1 \right]_{(X_{\theta_{j-1}}, t^{\frac{1}{4}} - K_{\theta_{j-1}})}^{(a, v)} \\ &\leq 25 r_2 \sup_{(a, v) \in [0,1] \times \mathbb{R}^+} \left\| \phi_{t, (a, v)} - \phi_\infty \right\|_1, \end{aligned} \quad (6.11)$$

where the first inequality follows from the definition of $c_{+,+}$ and (6.10) to which we apply Jensen's inequality over the measure determined by $|\phi_{t, (a, v)}(a', v') - \phi_\infty(a', v')| da' dv'$ and finally Hölder's inequality to pull the supremum outside the integral. The second inequality follows since $\mathbb{E}_{(x, k)} \left[\left| t^{-\frac{1}{4}} \int_0^\varsigma ds \frac{dV}{dx}(X_s) \right|^2 \right]$ is smaller than $25r_2$ for $(x, k) \in [0, 1] \times [0, t^{\frac{1}{20}}]$ by the same argument as in Part (1).

Finally by Proposition B.9, $\phi_{t, (a, v)}$ converges to ϕ_∞ in L^1 uniformly for $(a, v) \in [0, 1] \times [0, 2t^{\frac{1}{4}}]$ (which includes $[0, 1] \times [0, t^{\frac{1}{40}}]$) as $t \rightarrow \infty$.

Part (3):

Again we invoke Lemma 6.2, to work with the process conditioned to have jumps bounded by $t^{\frac{1}{40}}$. Let $\mathcal{F}_{\theta_{j-1}}$ be the σ -algebra of all information known up to the end of the last incursion θ_{j-1} , and j and m are related by (6.7). For a random process X_s , $0 \leq s \leq 1$, define $\|X_s\|_{p, \infty}$

for $p \geq 1$ as $\mathbb{E}[\sup_{0 \leq s \leq 1} |X_s|^p]^{\frac{1}{p}}$. By the triangle inequality and by Jensen's inequality for the first term on the right

$$\begin{aligned} \left\| \sum_{m=1}^{N_{\mathbf{s}_1}(st)} \left(Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - c_{\mathbf{s}_1, \mathbf{s}_2}(t) \right) \right\|_{1, \infty} &\leq \left\| \sum_{j=1}^{N_{\mathbf{s}_1}(st)} \left(Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}] \right) \right\|_{2, \infty} \\ &\quad + \left\| \sum_{j=1}^{N_{\mathbf{s}_1}(st)} \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}] - N_{\mathbf{s}_1}(st) c_{\mathbf{s}_1, \mathbf{s}_2}(t) \right\|_{1, \infty} \end{aligned} \quad (6.12)$$

Since the information of previous incursions is contained in $\mathcal{F}_{\theta_{j-1}}$, the sum of the differences $Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}]$ up to $m = N_{st}$ is a martingale. By Doob's inequality and Lemma A.1

$$\begin{aligned} \left\| \sum_{m=1}^{N_{\mathbf{s}_1}(st)} \left(Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}] \right) \right\|_{2, \infty} &\leq \mathbb{E} \left[\left| \sum_{m=1}^{N_{\mathbf{s}_1}(st)} Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}] \right|^2 \right]^{\frac{1}{2}} \\ &\leq \mathbb{E}[N_{\mathbf{s}_1}(st)]^{\frac{1}{2}} \sup_m \mathbb{E} \left[|Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}]|^2 \middle| m \leq N_{\mathbf{s}_1}(st) \right]^{\frac{1}{2}} \\ &\leq r_2^{\frac{1}{2}} t^{\frac{1}{8}} \sup_{m, \omega \in \mathcal{F}_{\sigma_j}} \mathbb{E} \left[|Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|^2 \middle| \mathcal{F}_{\sigma_j} \right]^{\frac{1}{2}} \leq 5r_2^{\frac{1}{2}} t^{\frac{1}{8}} \end{aligned} \quad (6.13)$$

where the last inequality uses Part (1), and the third inequality uses Part (2) of Lemma 6.1, and the following

$$\begin{aligned} &\mathbb{E} \left[|Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}]|^2 \middle| m \leq N_{\mathbf{s}_1}(st) \right]^{\frac{1}{2}} \\ &\leq \sup_{m, \omega \in \mathcal{F}_{\sigma_j}} \mathbb{E} \left[|Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}]|^2 \middle| \mathcal{F}_{\sigma_j} \right]^{\frac{1}{2}} \leq \sup_{m, \omega \in \mathcal{F}_{\sigma_j}} \mathbb{E} \left[|Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|^2 \middle| \mathcal{F}_{\sigma_j} \right]^{\frac{1}{2}}. \end{aligned} \quad (6.14)$$

Now we can work on the second term on the right of (6.12). By the triangle inequality and conditioning that $m \leq N_{\mathbf{s}_1}(st)$ for the terms in the sum as above,

$$\begin{aligned} &\mathbb{E} \left[\left| \sum_{m=1}^{N_{\mathbf{s}_1}(st)} \mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}] - N_{\mathbf{s}_1}(st) c_{\mathbf{s}_1, \mathbf{s}_2}(t) \right| \right] \\ &\leq \mathbb{E} \left[\sum_{m=1}^{N_{\mathbf{s}_1}(st)} \mathbb{E} \left[|\mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t)| \middle| m \leq N_{\mathbf{s}_1}(st) \right] \right] \end{aligned}$$

However, we will split the terms in the sum on the right-side into the two groups $m \in [N_{\mathbf{s}_1}(st - \varrho t) + 1, N_{\mathbf{s}_1}(st)]$ and $m \in [0, N_{\mathbf{s}_1}(st - \varrho t)]$ for some $0 < \varrho \ll 1$ and in particular $\varrho < s$. For $m \in [N_{\mathbf{s}_1}(st - \varrho t) + 1, N_{\mathbf{s}_1}(st)]$

$$\begin{aligned} &\mathbb{E} \left[\sum_{m=N_{\mathbf{s}_1}(st - \varrho t) + 1}^{N_{\mathbf{s}_1}(st)} \mathbb{E} \left[|\mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m) | \mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t)| \middle| m \leq N_{\mathbf{s}_1}(st) \right] \right] \\ &\leq 2\mathbb{E}[N_{\mathbf{s}_1}(st) - N_{\mathbf{s}_1}(st - \varrho t)] \sup_{\omega \in \mathcal{F}_{\sigma_j}} \mathbb{E}[|Y_{\mathbf{s}_1, \mathbf{s}_2}(m)| | \mathcal{F}_{\sigma_j}] \leq 5r_2^{\frac{1}{2}} \varrho^{\frac{1}{2}} t^{\frac{1}{4}}, \end{aligned}$$

where the first inequality follows since $\mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|\mathcal{F}_{\theta_{j-1}}]$ and $c_{\mathbf{s}_1, \mathbf{s}_2}(t)$ are convex combinations of values $\mathbb{E}[|Y_{\mathbf{s}_1, \mathbf{s}_2}(m)||\mathcal{F}_{\sigma_j}]$ for different $\omega \in \mathcal{F}_{\sigma_j}$. The second inequality employs Part (1) and then Part (2) of Lemma 6.1 for an interval of length ϱt .

For the sum of the terms with $m \in [0, N_{\mathbf{s}_1}(st - \varrho t)]$, we need to better understand the expressions

$$\mathbb{E}\left[\left|\mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|\mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t)\right| \middle| m \leq N_{\mathbf{s}_1}(st)\right], \quad (6.15)$$

and, in particular, how the information $m \leq N_{\mathbf{s}_1}(st)$ will change the expectation. If $m \leq N_{\mathbf{s}_1}(st)$, then it will already be known at time $\theta_{j-1} \leq (s - \varrho)t$, that $m - 1 \leq N_{\mathbf{s}_1}(st)$. However, it was shown in the beginning of the proof of Corollary B.9 that the probability of a jump into the region $[-t^{\frac{1}{4}}, t^{\frac{1}{4}}]$ (which is the beginning of an incursion) after starting in any point (x, k) with $|k| \in [2t^{\frac{1}{4}}, 4t^{\frac{1}{4}}]$ occurs with probability approaching one for $t \rightarrow \infty$. Since, we have assumed jumps bounded by $t^{\frac{1}{40}}$, the point $(X_{\theta_{j-1}}, K_{\theta_{j-1}})$ will have $|K_{\theta_{j-1}}| \leq [2t^{\frac{1}{4}}, 2t^{\frac{1}{4}} + t^{\frac{1}{40}}]$. Thus knowing $m \leq N_{\mathbf{s}_1}(st)$ will add little is known at time θ_j . With this consideration, we can give a crude upper bound for the expression (6.15) by doubling the unconditioned value of the expectation of $|Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|\mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t)|$:

$$\begin{aligned} \mathbb{E}\left[\left|\mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|\mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t)\right| \middle| N_{\mathbf{s}_1}(st) - N_{\mathbf{s}_1}(\theta_{j-1}) > 0\right] \\ < 2\mathbb{E}\left[\left|\mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|\mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t)\right|\right] \end{aligned} \quad (6.16)$$

where the inequality is due to the event $N_{\mathbf{s}_1}(st) - N_{\mathbf{s}_1}(\theta_{j-1}) > 0$ having probability close to one by Corollary B.9. Finally,

$$\begin{aligned} \mathbb{E}\left[t^{-\frac{1}{4}} \sum_{m=1}^{N_{\mathbf{s}_1}(st - \varrho t)} \mathbb{E}\left[\left|\mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|\mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t)\right|\right]\right] \\ < 2t^{-\frac{1}{4}}\mathbb{E}[N_{\mathbf{s}_1}(st - \varrho t)] \sup_m \mathbb{E}\left[\left|\mathbb{E}[Y_{\mathbf{s}_1, \mathbf{s}_2}(m)|\mathcal{F}_{\theta_{j-1}}] - c_{\mathbf{s}_1, \mathbf{s}_2}(t)\right|\right], \end{aligned}$$

which goes to zero by Part (2) of this proposition and by Part (2) of Lemma 6.1. $\square$

6.2 The torus reflection symmetry

Now we move on to results which are specific to having a reflection symmetry on the torus such that the potential $V(x)$ and jump densities $j_x(v)$ satisfy

$$V(x) = V(R(x)) \quad \text{and} \quad j_x(v) = j_{R(x)}(v).$$

A consequence of this symmetry along with the symmetric jump rates $j_a(v) = j_a(-v)$ between positive and negative momenta is that a specific phase space trajectory (x_s, k_s) , $s \in [0, t]$ from (x_0, k_0) to (x_t, k_t) will occur with the same density as a trajectory $(R(x_s), -k_s)$ from $(R(x_0), -k_0)$ to $(R(x_t), -k_t)$. This combines with the time reversal symmetry of the model to yield a “time reversal in momentum”.

Original trajectory (x_s, k_s)	Torus reflection ($R(x_s), -k_s$)
Time reversal symmetry ($x_{t-s}, -k_{t-s}$)	Time rev. with torus reflection ($R(x_{t-s}), k_{t-s}$)

The significance of the symmetries for the study of the drift is that $\frac{dV}{dx}(x) = -\frac{dV}{dx}(R(x_s))$. Thus for the trajectories (x_s, k_s) and $(R(x_{t-s}), k_{t-s})$ over the interval $[0, t]$,

$$\int_0^t ds \frac{dV}{dx}(x_s) = - \int_0^t ds \frac{dV}{dx}(R(x_{t-s})).$$

This will provide a basis for showing that there is no systematic bias for the contributions of the incursions.

Lemma 6.5. *Assume List 2.1 and List 2.2. For the momentum-capped dynamics $\frac{\rho_{+,-}(t)}{\rho_{-,+}(t)} - 1$ tends to zero.*

Proof. By the torus reflection symmetry of the dynamics, the probability density of going from $(a, t^{\frac{1}{4}} - v)$ to $(a', -2t^{\frac{1}{4}} - v')$ from the initial time to time θ is the same as the probability density of going from $(R(a), -t^{\frac{1}{4}} + v)$ to $(R(a'), 2t^{\frac{1}{4}} + v')$. Thus

$$\Pr_{(a, t^{\frac{1}{4}} - v)}[K_\theta < 0] = \Pr_{(R(a), -t^{\frac{1}{4}} + v)}[K_\theta > 0].$$

By the triangle inequality,

$$\left| \frac{\rho_{+,-}(t)}{\rho_{-,+}(t)} - 1 \right| \leq \left| \frac{\rho_{+,-}(t)}{\rho_{-,+}(t)} - \frac{\rho_{+,-}(t)}{\Pr_{(R(a), -t^{\frac{1}{4}} + v)}[K_\theta > 0]} \right| + \left| \frac{\rho_{+,-}(t)}{\Pr_{(a, t^{\frac{1}{4}} - v)}[K_\theta < 0]} - 1 \right| \quad (6.17)$$

By Proposition 6.3,

$$\left| \frac{\Pr_{(a, t^{\frac{1}{4}} - v)}[K_\theta < 0]}{\rho_{+,-}} - 1 \right|, \quad \left| \frac{\Pr_{(a, -t^{\frac{1}{4}} + v)}[K_\theta > 0]}{\rho_{-,+}} - 1 \right| \longrightarrow 0,$$

for all $(a, v) \in [0, 1] \times [0, t^{\frac{1}{40}}]$. Thus (6.17) goes to zero. $\square$

The following lemma constructs a specific joint distribution $\phi_t^*(a, v; a', v')$ for the first entrance coordinates (a, v) and last exit coordinates (a', v') for the set $S = \{(x, k) \in [0, 1] \times \mathbb{R} \mid |k| \leq t^{\frac{1}{4}}\}$ for trajectories conditioned to begin with $s_1 K_0 > t^{\frac{1}{4}}$ and to end with $s_2 K_\theta > t^{\frac{1}{4}}$. The symmetry (6.18) will play a key role in the proof of Proposition 6.7.

Lemma 6.6 (Equilibrium first-entrance/last-exit distribution). *Assume List 2.1 and List 2.2. Consider the dynamics conditioned to have jumps capped by $t^{\frac{1}{40}}$. For large enough t , there exists a unique joint density $\phi_t^*(a, v; a', v')$ with support in $([0, 1] \times [0, t^{\frac{1}{40}}])^2$ such that the marginals*

$$\phi_{\mathcal{F},t}^*(a, v) = \int_{[0,1] \times \mathbb{R}^+} da' dv' \phi_t^*(a, v; a', v') \quad \text{and} \quad \phi_{\mathcal{L},t}^*(a', v') = \int_{[0,1] \times \mathbb{R}^+} da dv \phi_t^*(a, v; a', v'),$$

satisfy the relations

$$\begin{aligned}\phi_{\mathcal{L},t}^*(a', v') &= \int_{[0,1] \times \mathbb{R}^+} da dv \phi_{\mathcal{F},t}^*(a, v) \Pr_{(a, s_1 t^{\frac{1}{4}} - s_1 v)}[(X_\varsigma, s_2 K_\varsigma) = (a', t^{\frac{1}{4}} - v') \mid s_2 K_\varsigma > 0] \\ \phi_{\mathcal{F},t}^*(a', v') &= \int_{[0,1] \times \mathbb{R}^+} da dv \phi_{\mathcal{L},t}^*(a, v) \Pr_{(a, s_1 t^{\frac{1}{4}} - s_1 v)}^{(R)}[(X_\varsigma, s_2 K_\varsigma) = (a', t^{\frac{1}{4}} - v') \mid s_2 K_\varsigma > 0,]\end{aligned}$$

where $\Pr_{(x,k)}^{(R)}$ refers to the law of the time-reversed dynamics starting from the point $(x, k) \in [0, 1] \times \mathbb{R}$.

Moreover, ϕ_t^* has the symmetry

$$\phi_t^*(a, v; a', v') = \phi_t^*(R(a'), v'; R(a), v). \quad (6.18)$$

Proof. Set $s_1 = s_2 = +$. We pick a number in $(1, 2)$, say, $\frac{3}{2}$. In the event that $K_\sigma > 0$, define ς' to be the last time the particle has a jump with momentum $< \frac{3}{2} t^{\frac{1}{4}}$ before it continues on to reach $2 t^{\frac{1}{2}}$ (at time θ),

$$\varsigma' = \inf\{s \in [0, \theta], M_s - M_{s-} \neq 0 \mid \inf_{s < r \leq \theta} |K_r| > \frac{3}{2} t^{\frac{1}{4}}\}. \quad (6.19)$$

Clearly $\varsigma' > \varsigma$.

Consider the two maps $\Psi^F, \Psi^R : L^1([0, 1] \times \mathbb{R}^+)$

$$\begin{aligned}\Psi^F(\varphi)(a, v) &= \int da' dv' \varphi(a, v) \Pr_{(a, \frac{3}{2} t^{\frac{1}{4}} - v)}[(X_{\varsigma'}, K_{\varsigma'}) = (a', \frac{3}{2} t^{\frac{1}{4}} - v') \mid K_\varsigma > 0, \inf_{0 \leq r \leq \theta} K_r < t^{\frac{1}{4}}] \\ \Psi^R(\varphi)(a, v) &= \int da' dv' \varphi(a, v) \Pr_{(a, \frac{3}{2} t^{\frac{1}{4}} - v)}^{(R)}[(X_\varsigma, K_\varsigma) = (a', \frac{3}{2} t^{\frac{1}{4}} - v') \mid K_\varsigma > 0, \inf_{0 \leq r \leq \theta} K_r < t^{\frac{1}{4}}],\end{aligned}$$

where $\Pr_{(x,k)}^{(R)}$ refers to the statistics for the time-reversed Markov dynamics starting from the point (x, k) .

Ψ^F and Ψ^R send probability densities to probability densities, and for large enough t , we claim that Ψ^F, Ψ^R are contractive on differences of densities. Consider the hitting time $\tau = \inf\{s \in [0, \theta], M_s - M_{s-} \neq 0 \mid K_s \notin S\}$ for the set $S = \{(x, k) \in [0, 1] \times \mathbb{R} \mid |k - \frac{3}{2} t^{\frac{1}{4}}| < t^{\frac{1}{20}}\}$.

Since $\tau < \varsigma'$, by the Markov property,

$$\begin{aligned}\Psi^F(\varphi)(a, v) &= \int_{[0,1] \times \mathbb{R}^+} da' dv' \varphi(a', v') \\ &\quad \times \int_{[0,1] \times \mathbb{R}^+} dx dk (\psi_{\uparrow,t}^{(a',v')}(x, k - \frac{3}{2} t^{\frac{1}{4}} - t^{\frac{1}{20}}) + \psi_{\downarrow,t}^{(a',v')}(x, -k + \frac{3}{2} t^{\frac{1}{4}} - t^{\frac{1}{20}})) \\ &\quad \times \Pr_{(x,k)}[(X_{\varsigma'}, K_{\varsigma'}) = (a, \frac{3}{2} t^{\frac{1}{4}} - v) \mid K_\varsigma > 0, \inf_{0 \leq r \leq \theta} K_r < t^{\frac{1}{4}}] \quad (6.20)\end{aligned}$$

$\psi_{\uparrow,t}^{(a',v')}$ and $\psi_{\downarrow,t}^{(a',v')}$ are the boundary crossing densities for the set S for paths starting from $(a', \frac{3}{2} t^{\frac{1}{4}} - v')$ and conditioned to reach below $t^{\frac{1}{4}}$ before going above $t^{\frac{1}{4}}$. Using (6.20), for two probability densities φ_1, φ_2 ,

$$\begin{aligned}\|\Psi^F(\varphi_1 - \varphi_2)\|_1 &\leq \|\varphi_1 - \varphi_2\|_1 \\ &\quad \times \sup_{a, a' \in [0,1], v, v' \in [0, t^{\frac{1}{40}}]} (\|\psi_{\uparrow,t}^{(a,v)} - \psi_{\uparrow,t}^{(a',v')}\|_1 + \|\psi_{\downarrow,t}^{(a,v)} - \psi_{\downarrow,t}^{(a',v')}\|_1). \quad (6.21)\end{aligned}$$

However, by Proposition B.8 we have that

$$\sup_{a \in [0,1], v, v' \in [0, t^{\frac{1}{40}}]} \|\psi_{\uparrow, t}^{(a, v)} - \frac{1}{2}\phi_{\infty}\|_1 \longrightarrow 0 \quad \text{and} \quad \sup_{a \in [0,1], v \in [0, t^{\frac{1}{40}}]} \|\phi_{\downarrow, t}^{(a, v)} - \frac{1}{2}\phi_{\infty}\|_1 \longrightarrow 0.$$

Thus for large enough t there exists a constant $0 < \lambda < 1$ such that

$$\|\Psi^F(\varphi_1 - \varphi_2)\|_1 \leq \lambda \|\varphi_1 - \varphi_2\|_1,$$

for any two probability densities $\varphi_1, \varphi_2 \in L^1([0, 1] \times \mathbb{R}^+)$. For large enough t , the constant $\lambda > 0$ can be made arbitrarily small. By symmetry, the same proof holds for Ψ^R .

For t large enough for the strict contractive property above, we can now construct special densities by defining the limits

$$\pi_t^F = \lim_{n \rightarrow \infty} (\Psi^R \Psi^F)^n(P) \quad \text{and} \quad \pi_t^R = \lim_{n \rightarrow \infty} (\Psi^F \Psi^R)^n(P),$$

where the limit is independent of the probability density P . We have constructed equilibrium states in which we can do computations using the Markov property for both the original dynamics and the time-reversed dynamics.

We thus define the first entrance distribution

$$\phi_{\mathcal{F}, t}^*(a, v) = \int_{[0,1] \times \mathbb{R}^+} da' dv' \pi_t^F(a', v') \Pr_{(a, t^{\frac{1}{4}} - v)}[(X_{\tau}, K_{\tau}) = (a', t^{\frac{1}{4}} - v') \mid K_{\theta} > 0 \inf_{0 \leq r \leq \theta} K_r < t^{\frac{1}{4}}], \quad (6.22)$$

and finally $\phi_t^*(a, v; a', v')$ as the product

$$\phi_t^*(a, v; a', v') = \phi_{\mathcal{F}, t}^*(a, v) \Pr_{(a, t^{\frac{1}{4}} - v)}[(X_{\varsigma}, K_{\varsigma}) = (a', t^{\frac{1}{4}} - v') \mid K_{\varsigma} > 0].$$

The first relation in the statement of the lemma, which determines $\phi_{\mathcal{L}, t}^*(a, v)$ with $\phi_{\mathcal{F}, t}^*(a, v)$ using the forward dynamics, follows immediately from the definition of ϕ_t^* . In the above formula, $\phi_{\mathcal{L}, t}^*(a', v')$ is determined as the last exit time of $S = \{(x, k) \in [0, 1] \times \mathbb{R} \mid |k| \leq t^{\frac{1}{4}}\}$. However, starting from the points $(q, \frac{3}{2}t^{\frac{1}{4}} - p)$ with distribution $\pi_t^R(q, p)$ in the time-reversed dynamics, then $\phi_{\mathcal{L}, t}^*(a, v)$ is the first entrance distribution for set S . The second relation then follows from the Markov property for the time-reversed dynamics

$$\phi_{\mathcal{F}, t}^*(a, v) = \int da' dv' \phi_{\mathcal{L}, t}^*(a', v') \Pr_{(a, t^{\frac{1}{4}} - v)}^{(R)}[(X_{\varsigma}, K_{\varsigma}) = (a', t^{\frac{1}{4}} - v') \mid K_{\varsigma} > 0]. \quad (6.23)$$

□

Proposition 6.7 (Antisymmetry of constants). *Assume List 2.1 and List 2.2. The constants $c_{+, +}(t)$, $c_{-, -}(t)$, $c_{+, -}(t) + c_{-, +}(t)$ tend to zero for large times.*

Proof. Let us fix $\mathbf{s}_1 = \mathbf{s}_2 = +$. By Lemma 6.2, we can take the dynamics conditioned to make jumps capped by $t^{\frac{1}{40}}$.

Define the functional $\Lambda : L^1([0, 1] \times \mathbb{R}^+) \rightarrow \mathbb{R}$,

$$\Lambda(\varphi) = \int_{[0,1] \times [0, t^{\frac{1}{40}}]} da dv \varphi(a, v) \mathbb{E}_{(a, t^{\frac{1}{4}} - v)}[t^{-\frac{1}{4}} \int_0^{\varsigma} ds \frac{dV}{dx}(X_s) \chi(K_{\varsigma} > 0)]. \quad (6.24)$$

By our comments above $c_{+,+} \approx \Lambda(\phi_\infty)$, and when $\varphi = \phi_{\mathcal{F},t}^*$ then

$$\Lambda(\phi_{\mathcal{F},t}^*) = \Pr[K_\theta > 0] \int da dv da' dv' \phi_t^*(a, v; a', v') \mathbb{E}_{(a, t^{\frac{1}{4}}-v)}^{(a', t^{\frac{1}{4}}-v')} \left[t^{-\frac{1}{4}} \int_0^\varsigma ds \frac{dV}{dx}(X_s) \right], \quad (6.25)$$

where $\mathbb{E}_{(x,k)}^{(x',k')}$ is the expectation conditioned on the trajectories that begin at (x, k) and have a last exit (x', k') at the time ς .

Note the anti-symmetry

$$\mathbb{E}_{(a, t^{\frac{1}{4}}-v)}^{(a', t^{\frac{1}{4}}-v')} \left[t^{-\frac{1}{4}} \int_0^\varsigma ds \frac{dV}{dx}(X_s) \right] = -\mathbb{E}_{(R(a'), t^{\frac{1}{4}}-v')}^{(R(a), t^{\frac{1}{4}}-v)} \left[t^{-\frac{1}{4}} \int_0^\varsigma ds \frac{dV}{dx}(X_s) \right], \quad (6.26)$$

which is due to the “time-reversal in momentum” mentioned at the beginning of the section in which for every trajectory (x_t, k_t) on the interval $[0, t]$, there is a backwards trajectory with a torus-reflected position $(R(x_{t-s}), k_{t-s})$ which occurs in the forward dynamics with the same “probability” as a fraction of the trajectories that begin at (x_0, k_0) and (x_t, k_t) respectively. Even though the final time ς is not deterministic, the two trajectories are still weighted equally in the expectations (6.26). This follows since ς is a hitting time for the time-reversed Markov process (for when the momentum first jumps below $t^{\frac{1}{4}}$). Due to (6.26) and the symmetry (6.18) of $\phi_t^*(a, v; a', v')$, it follows that (6.25) is zero.

We now focus on showing that due to a dynamical loss of memory of the initial conditions over the time interval $[0, \sigma]$, the values of $\Lambda(\varphi)$ for any φ with support in $[0, 1] \times [0, t^{\frac{1}{40}}]$ are close. In particular, $\Lambda(\varphi)$ for $\varphi = \bar{\phi}_\infty = \phi_\infty \chi(|v| \leq t^{\frac{1}{40}})$ or $\varphi = \phi_{\mathcal{F},t}^*$ are close, which would prove the result.

Define τ to be the exit time for the set $S = \{(x, k) \in [0, 1] \times \mathbb{R} \mid |k - t^{\frac{1}{4}}| < t^{\frac{1}{20}}\}$. Thus for a probability density φ ,

$$\begin{aligned} \Lambda(\varphi) = & \int_{[0,1] \times \mathbb{R}^+} da dv \varphi(a, v) \mathbb{E}_{(a, t^{\frac{1}{4}}-v)} \left[t^{-\frac{1}{4}} \int_0^{\tau \wedge \sigma} ds \frac{dV}{dx}(X_s) \chi(K_\varsigma > 0) \right] \\ & + \int dx dk \left(\phi_{\uparrow, t}^\varphi(x, k - t^{\frac{1}{4}} - t^{\frac{1}{20}}) + \phi_{\downarrow, t}^\varphi(x, -k + t^{\frac{1}{4}} - t^{\frac{1}{20}}) \right) \\ & \times \mathbb{E}_{(x, k)} \left[t^{-\frac{1}{4}} \int_0^\sigma ds \frac{dV}{dx}(X_s) \chi(K_\varsigma > 0) \right]. \end{aligned} \quad (6.27)$$

where $\phi_{\uparrow, t}^\varphi = \int dq dp \varphi(q, p) \phi_{\uparrow, t}^{(q, p)}$ and an analogous definition for $\phi_{\downarrow, t}^\varphi$.

We argue that the first term on the right-side of (6.27) tends to zero for large t . Since the momentum is greater than $\frac{1}{2}t^{\frac{1}{4}}$ up to time τ , then by Lemma 5.1

$$\left| \mathbb{E}_{(a, t^{\frac{1}{4}}-v)} \left[t^{-\frac{1}{4}} \int_0^{\tau \wedge \sigma} ds \frac{dV}{dx}(X_s) \right] \right| \leq \frac{4\bar{V}}{t^{\frac{1}{4}}} \mathbb{E}_{(a, t^{\frac{1}{4}}-v)} [N_\tau],$$

where N_τ is the number of momentum jumps up to time τ . In the proof of Proposition B.6, it was shown that $\mathbb{E}_{(x, k)} [N] = O(t^{\frac{1}{10}})$. Thus the drift up to time $\tau \wedge \sigma$ vanishes for large t vanishes as $O(t^{-\frac{3}{20}})$.

Thus

$$\begin{aligned} |\Lambda(\phi) - \Lambda(\phi_{\mathcal{F},t}^*)| \leq & O(t^{-\frac{3}{20}}) + \left(\|\phi_{\uparrow, t}^\varphi|_{\varphi=\bar{\phi}_\infty} - \phi_{\uparrow, t}^\varphi|_{\varphi=\phi_{\mathcal{F},t}^*}\|_1 + \|\phi_{\downarrow, t}^\varphi|_{\varphi=\bar{\phi}_\infty} - \phi_{\downarrow, t}^\varphi|_{\varphi=\phi_{\mathcal{F},t}^*}\|_1 \right) \\ & \times \sup_{(x, k)} \mathbb{E}_{(x, k)} \left[\left| t^{-\frac{1}{4}} \int_0^\sigma ds \frac{dV}{dx}(X_s) \chi(K_\varsigma > 0) \right| \right]. \end{aligned}$$

However, by Proposition B.7, $\phi_{\uparrow,t}^\varphi, \phi_{\downarrow,t}^\varphi \rightarrow \varphi_\infty$, since by definition $\phi_{\uparrow,t}^\varphi$ is a convex combination of $\phi_{\uparrow,t}^{(q,p)}$ for (a,p) with $p \in [0, t^{\frac{1}{40}}]$.

$\phi_{\uparrow,t}^\varphi|_{\varphi=\phi_\infty}, \phi_{\uparrow,t}^\varphi|_{\varphi=\phi_{\mathcal{F},t}^*}, \phi_{\downarrow,t}^\varphi|_{\varphi=\phi_\infty}$, and $\phi_{\downarrow,t}^\varphi|_{\varphi=\phi_{\mathcal{F},t}^*}$ tend to ϕ_∞ in L^1 . Moreover, by the same argument as for Part (1) of Proposition 6.4,

$$\sup_{(x,k)} \left| \mathbb{E}_{(x,k)} \left[\left| t^{-\frac{1}{4}} \int_0^\sigma ds \frac{dV}{dx}(X_s) \right|^2 \chi(K_\theta > 0) \right] \right|^{\frac{1}{2}} \leq 5.$$

We then have that $|\Lambda(\phi) - \Lambda(\phi_{\mathcal{F},t}^*(a, v))|$ converges to zero for $t \rightarrow \infty$ which proves the result. That shows the $c_{+,+}$ case. The other cases for $c_{-,-}$, and $c_{+,-} + c_{-,+}$ are similar. $\square$

Theorem 6.8. *Assume List 2.1 and List 2.2. In the limit $t \rightarrow \infty$,*

$$\mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| t^{-\frac{1}{2}} \int_0^{st} dr \frac{dV}{dx}(X_r) \right| \right] \rightarrow 0. \quad (6.28)$$

Proof. Our basic idea is to break the integral $t^{-\frac{1}{2}} \int_0^{st} dr \frac{dV}{dx}(X_r)$ into parts where corresponding to where $|K_r|$ is high and low energy respectively. The low energy parts are controlled by our study of the random variables $Y_n = \int_{\sigma_n}^{\varsigma_n} dr \frac{dV}{dx}(X_r)$ and the high energy parts will be controlled by Lemma 5.3.

$$\begin{aligned} t^{-\frac{1}{2}} \int_0^{st} dr \frac{dV}{dx}(X_r) &= t^{-\frac{1}{2}} \int_0^{\theta_0} dr \frac{dV}{dx}(X_r) + t^{-\frac{1}{2}} \int_0^{st} dr \frac{dV}{dx}(X_r) \chi(|K_{\tau(r)}| > t^{\frac{1}{4}}) \\ &\quad + t^{-\frac{1}{4}} \sum_{n=1}^{N_Y(st)} Y_n - \int_{\sigma_n}^{\varsigma_n} dr \frac{dV}{dx}(X_r) \chi(|K_{\tau(r)}| > t^{\frac{1}{4}}) \\ &\quad - \chi(\exists(j) : st \in [\sigma_j, \varsigma_j]) \int_{st}^{\theta_{N_Y(st)}} dr \frac{dV}{dx}(X_r) \end{aligned} \quad (6.29)$$

where $\tau(r)$ the next to last jump time before time r as in Lemma 5.3, and the last term corresponds removing an overlap due to a last incomplete incursion which begins before

For the first term on the right side of (6.29), an argument analogous to Part (1) of Proposition 6.4 gives the bound,

$$\mathbb{E} \left[\left| \int_0^{\theta_0} dr \frac{dV}{dx}(X_r) \right|^2 \right]^{\frac{1}{2}} \leq 5 t^{\frac{1}{4}}, \quad (6.30)$$

and thus that term is negligible. In this case, the end time γ is a hitting time, which makes the argument easier. The last term has the same bound. By Lemma 5.3,

$$\mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| t^{-\frac{1}{2}} \int_0^{st} dr \frac{dV}{dx}(X_r) \chi(|K_{\tau(r)}| > t^{\frac{1}{4}}) \right|^2 \right]^{\frac{1}{2}}$$

converges to zero. The same argument as in the proof of Lemma 5.3 shows that

$$\mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| t^{-\frac{1}{2}} \sum_{n=1}^{N_Y(st)} \int_{\sigma_n}^{\varsigma_n} dr \frac{dV}{dx}(X_r) \chi(|K_{\tau(r)}| > t^{\frac{1}{4}}) \right|^2 \right]^{\frac{1}{2}} \rightarrow 0,$$

since it includes even less terms.

We are left with the sequence $t^{-\frac{1}{4}} \sum_{n=1}^{N_Y(st)-1} Y_n$, which we can write as

$$t^{-\frac{1}{4}} \sum_{m=1}^{N_+(st)} Y_{+,+}(m) + Y_{+,-}(m) + t^{-\frac{1}{4}} \sum_{m=1}^{N_-(st)} Y_{-,-}(m) + Y_{-,+}(m).$$

By Part (4) of Proposition 6.4 these sums can be approximated by $t^{-\frac{1}{4}} N_{\mathbf{s}_1}(st) c_{\mathbf{s}_1, \mathbf{s}_2}(t)$:

$$\mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| t^{-\frac{1}{4}} \sum_{j=1}^{N_{\mathbf{s}_1}(st)} Y_{\mathbf{s}_1, \mathbf{s}_2}(m) - t^{-\frac{1}{4}} N_{\mathbf{s}_1}(st) c_{\mathbf{s}_1, \mathbf{s}_2}(t) \right| \right] \rightarrow 0.$$

For the sequences with $\mathbf{s}_1 = \mathbf{s}_2 = \mathbf{s}$

$$\mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| t^{-\frac{1}{4}} N_{\mathbf{s}, \mathbf{s}}(st) c_{\mathbf{s}, \mathbf{s}_1}(t) \right| \right] = \mathbb{E} \left[\left| t^{-\frac{1}{4}} N_{\mathbf{s}, \mathbf{s}}(t) c_{\mathbf{s}, \mathbf{s}}(t) \right| \right] \leq r_2^{\frac{1}{2}} |c_{\mathbf{s}, \mathbf{s}}(t)|,$$

where the inequality uses Part (2) of Lemma 6.1. By Proposition 6.7, $c_{\mathbf{s}, \mathbf{s}}(t)$ converges to zero.

The cases of $(\mathbf{s}_1, \mathbf{s}_2) = (+, -)$ and $(\mathbf{s}_1, \mathbf{s}_2) = (-, +)$ must be treated together. We will take a step backward and approximate $t^{-\frac{1}{4}} N_{\mathbf{s}_1}(st) c_{\mathbf{s}_1, \mathbf{s}_2}(t)$ with $t^{-\frac{1}{4}} (\rho_{\mathbf{s}_1, \mathbf{s}_2}(t))^{-1} \sum_{n=1}^{N_{\mathbf{s}_1}(st)} c_{\mathbf{s}_1, \mathbf{s}_2}(t) \chi(\mathbf{s}_2 K_{\zeta_j} > 0)$. By the triangle inequality

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| t^{-\frac{1}{4}} (\rho_{\mathbf{s}_1, \mathbf{s}_2}(t))^{-1} \sum_{n=1}^{N_{\mathbf{s}_1}(st)} c_{\mathbf{s}_1, \mathbf{s}_2}(t) \chi(\mathbf{s}_2 K_{\zeta_j} > 0) - t^{-\frac{1}{4}} N_{\mathbf{s}_1}(st) c_{\mathbf{s}_1, \mathbf{s}_2}(t) \right| \right] \\ & \leq t^{-\frac{1}{4}} (\rho_{\mathbf{s}_1, \mathbf{s}_2}(t))^{-1} |c_{\mathbf{s}_1, \mathbf{s}_2}(t)| \mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| \sum_{n=1}^{N_{\mathbf{s}_1}(st)} \chi(\mathbf{s}_2 K_{\zeta_j} > 0) - \Pr[\mathbf{s}_2 K_{\sigma_j} > 0] \right| \right] \\ & \quad + t^{-\frac{1}{4}} (\rho_{\mathbf{s}_1, \mathbf{s}_2}(t))^{-1} c_{\mathbf{s}_1, \mathbf{s}_2}(t) \mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| \sum_{n=1}^{N_{\mathbf{s}_1}(st)} \Pr[\mathbf{s}_2 K_{\zeta_j} > 0] - \rho_{\mathbf{s}_1, \mathbf{s}_2}(t) \right| \right]. \quad (6.31) \end{aligned}$$

The sum $\sum_{n=1}^{N_{\mathbf{s}_1}(st)} \chi(\mathbf{s}_2 K_{\zeta_j} > 0) - \Pr[\mathbf{s}_2 K_{\zeta_j} > 0]$ is a martingale, so by Doob's maximal inequality and by Lemma A.1

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| \sum_{n=1}^{N_{\mathbf{s}_1}(st)} \chi(\mathbf{s}_2 K_{\zeta_j} > 0) - \Pr[\mathbf{s}_2 K_{\zeta_j} > 0] \right|^2 \right]^{\frac{1}{2}} \leq 2 \mathbb{E} \left[\left| \sum_{n=1}^{N_{\mathbf{s}_1}(st)} \chi(\mathbf{s}_2 K_{\zeta_j} > 0) - \Pr[\mathbf{s}_2 K_{\zeta_j} > 0] \right|^2 \right]^{\frac{1}{2}} \\ & \leq \sup_j \mathbb{E} \left[(\chi(\mathbf{s}_2 K_{\zeta_j} > 0) - \Pr[\mathbf{s}_2 K_{\zeta_j} > 0])^2 | j \leq N_{\mathbf{s}_1}(st) \right]^{\frac{1}{2}} \mathbb{E} [N_{\mathbf{s}_1}(t)]^{\frac{1}{2}} \\ & \leq \sup_{j, s} \left| \Pr[\mathbf{s}_2 K_{\zeta_j} > 0 | j \leq N_{\mathbf{s}_1}(st)] - \Pr[\mathbf{s}_2 K_{\zeta_j} > 0] \right| r_2 t^{\frac{1}{8}}, \quad (6.32) \end{aligned}$$

where the last inequality follows since $N_{\mathbf{s}_1}(t) < N_Y(t)$ and from Part (2) of Proposition 6.1 and by an explicit calculation for the expectation of the indicator in the variance-type formula. Since the factor of $t^{-\frac{1}{4}}$ in (6.31) over-powers the factor $t^{\frac{1}{8}}$, the only worry is $(\rho_{\mathbf{s}_1, \mathbf{s}_2}(t))^{-1}$ going to zero. Since the event $j \leq N_{\mathbf{s}_1}(st)$ is adapted to the information known up to time σ_j ,

$$\begin{aligned} & (\rho_{\mathbf{s}_1, \mathbf{s}_2}(t))^{-1} \sup_j \left| \Pr[\mathbf{s}_2 K_{\zeta_j} > 0 | j \leq N_{\mathbf{s}_1}(st)] - \Pr[\mathbf{s}_2 K_{\zeta_j} > 0] \right| \\ & \leq \sup_{j, \omega \in \mathcal{F}_{\sigma_j}} \left| \frac{\Pr[\mathbf{s}_2 K_{\zeta_j} > 0 | \mathcal{F}_{\sigma_j}]}{\rho_{\mathbf{s}_1, \mathbf{s}_2}(t)} - \frac{\Pr[\mathbf{s}_2 K_{\zeta_j} > 0]}{\rho_{\mathbf{s}_1, \mathbf{s}_2}(t)} \right|. \quad (6.33) \end{aligned}$$

By adding and subtracting 1 in the expression on the right-side, then by two applications of Lemma 6.3, which is permitted by our assumption on the boundedness of the jumps, shows that the above goes to zero. For the application of Lemma 6.3, note that by the definitions θ_j and ς_j that $\mathbf{s}_2 K_{\varsigma_j} > 0$ is equivalent to $\mathbf{s}_2 K_{\theta_j} > 0$. Also note that $\Pr[\mathbf{s}_2 K_{\varsigma_j} > 0]$ is a convex combination of the probabilities $\Pr[\mathbf{s}_2 K_{\varsigma_j} > 0 | (X_{\sigma_j}, K_{\sigma_j})]$. Due to the decay of $t^{-\frac{1}{8}}$, we had only needed this term to be bounded, but we apply these principles again below.

For the second term in (6.31),

$$t^{-\frac{1}{4}} \mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| \sum_{n=1}^{N_{\mathbf{s}_1}(st)} \frac{\Pr[\mathbf{s}_2 K_{\varsigma_j} > 0]}{\rho_{\mathbf{s}_1, \mathbf{s}_2}(t)} - 1 \right| \right] \leq t^{-\frac{1}{4}} \sup_j \left| \frac{\Pr[\mathbf{s}_2 K_{\varsigma_j} > 0]}{\rho_{\mathbf{s}_1, \mathbf{s}_2}(t)} - 1 \right| \mathbb{E}[N_Y].$$

We apply Part (2) of Lemma 6.1 to show that $t^{-\frac{1}{4}} \mathbb{E}[N_Y] \leq r_2^{\frac{1}{2}}$ is bounded and Lemma 6.3 to show that the ratio of probabilities converges to one.

Now we just need to bound

$$t^{-\frac{1}{4}} (\rho_{+,-}(t))^{-1} \sum_{n=1}^{N_+(st)} c_{+,-}(t) \chi(K_{\varsigma_j} < 0) + t^{-\frac{1}{4}} (\rho_{-,(+)}(t))^{-1} \sum_{n=1}^{N_-(st)} c_{-,+}(t) \chi(K_{\varsigma_j} > 0). \quad (6.34)$$

Using Lemma 6.5 and the same techniques above, we can replace $\rho_{-,(+)}(t)$ by $\rho_{+,-}(t)$. More critically, since the number of up-crossings from below $-2t^{\frac{1}{4}}$ to above $2t^{\frac{1}{4}}$ can differ by at most one from the number of down-crossings from above $2t^{\frac{1}{4}}$ to below $-2t^{\frac{1}{4}}$,

$$\begin{aligned} & (\rho_{+,-}(t))^{-1} \mathbb{E} \left[\sup_{0 \leq s \leq 1} \left| t^{-\frac{1}{4}} \sum_{n=1}^{N_+(st)} c_{+,-}(t) \chi(K_{\varsigma_j} < 0) + t^{-\frac{1}{4}} \sum_{n=1}^{N_-(st)} c_{-,+}(t) \chi(K_{\varsigma_j} > 0) \right| \right] \\ & < \sup_j \left(\frac{\Pr[K_{\varsigma_j} < 0]}{\rho_{+,-}(t)} + \frac{\Pr[K_{\varsigma_j} > 0]}{\rho_{+,-}(t)} \right) \left(t^{-\frac{1}{4}} |c_{+,-}(t) + c_{-,+}(t)| \mathbb{E}[N_Y(t)] + c_{+,-}(t) \wedge c_{-,+}(t) \right) \end{aligned}$$

As above $\mathbb{E}[N_Y(t)] = O(t^{\frac{1}{4}})$, and by Lemmas 6.7 and 6.3

$$\sup_j \left(\frac{\Pr[K_{\varsigma_j} > 0]}{\rho_{+,-}(t)} + \frac{\Pr[K_{\varsigma_j} < 0]}{\rho_{+,-}(t)} \right) < 4.$$

Finally, $|c_{+,-}(t) + c_{-,+}(t)|$ converges to zero by Lemma 6.7, which finishes the proof. $\square$

7 Proof of main results

Proof of Theorem 3.1. Denote the initial distribution $P_0(x, k)$ as μ . We wish to apply the martingale central limit theorem. The Lindberg condition follows easily, since jumps occur at Poisson rate $\mathcal{R}$ and have finite fourth moments by (II) of List 2.1. Weak convergence with respect to the uniform metric then follows by convergence in probability of the predictable quadratic variation $t^{-1} \langle M \rangle_{st}$ to σs for every s . Without losing generality, we take $s = 1$ and show $\mathbb{E}_\mu[|t^{-1} \langle M \rangle_t - \sigma|] \rightarrow 0$.

Define $S_{\epsilon, \delta}^{(t)}$ to be the event that $\int_0^1 ds \chi(t^{-\frac{1}{2}} |K_{st}| > \epsilon) > 1 - \delta$. The same statement as in Lemma 3.4 holds for $t^{-\frac{1}{2}} E_{st}$ replaced by $t^{-\frac{1}{2}} |K_{st}|$, since $E_r^{\frac{1}{2}} \approx 2^{-\frac{1}{2}} |K_r|$ when $|K_r| \gg 1$.

Thus for some C , $\Pr[S_{\epsilon,\delta}^{(t)}] \geq 1 - C \frac{\sqrt{r_2}}{r_1} \frac{\epsilon}{\delta}$ for large enough t . It will be convenient to observe that $\mathcal{R} \int_0^r ds \chi(t^{-\frac{1}{2}}|K_r| \leq \epsilon)$ is close to the number $n_{r,\epsilon}$ of Poisson times t_m such that $|K_{t_m}^-| \leq \epsilon t^{\frac{1}{2}}$ up to time $r \leq t$. This is a law of large numbers following from the difference $n_{r,\epsilon} - \mathcal{R} \int_0^r ds \chi(t^{-\frac{1}{2}}|K_s| \leq \epsilon)$ being a martingale with

$$\mathbb{E}_\mu \left[\left(t^{-1} n_{t,\epsilon} - \mathcal{R} \int_0^1 ds \chi(t^{-\frac{1}{2}}|K_{st}| \leq \epsilon) \right)^2 \right]^{\frac{1}{2}} \leq t^{-\frac{1}{2}} \mathcal{R}. \quad (7.1)$$

Using conditional expectations and the triangle inequality over a telescoping sum determined by the Poisson times t_m , $m = 1, \dots, \mathcal{N}_t$, we get the first inequality below.

$$\begin{aligned} \mathbb{E}_\mu \left[\left| \frac{\langle M \rangle_t}{t} - \sigma \right| \right] &\leq \mathbb{E}_\mu \left[\frac{1}{t} \sum_{m=0}^{\mathcal{N}_t} \mathbb{E}_{(X_{t_m}, K_{t_m})} \left[\left| \langle M \rangle_{t_{m+1}} - \langle M \rangle_{t_m} - \sigma(t_{m+1} - t_m) \right| \right] \right] \\ &< 2^{\frac{1}{2}} r_2 (1 - \Pr[S_{\epsilon,\delta}^{(t)}])^{\frac{1}{2}} + \mathbb{E}_\mu \left[\chi(S_{\epsilon,\delta}^{(t)}) \frac{1}{t} \sum_{m=0}^{\mathcal{N}_t} \mathbb{E}_{(X_{t_m}, K_{t_m})} \left[\left| \langle M \rangle_{t_{m+1}} - \langle M \rangle_{t_m} - \sigma(t_{m+1} - t_m) \right| \right] \right]. \end{aligned} \quad (7.2)$$

The second inequality follows by using the brute upperbound for the complement of the event $S_{\epsilon,\delta}$:

$$\frac{1}{t} \sum_{m=0}^{\mathcal{N}_t} \mathbb{E}_{(X_{t_m}, K_{t_m})} \left[\left| \langle M \rangle_{t_{m+1}} - \langle M \rangle_{t_m} - \sigma(t_{m+1} - t_m) \right| \right] < r_2 \frac{\mathcal{N}_t}{t \mathcal{R}}, \quad (7.3)$$

which holds since $\langle M \rangle_r$ always increases at rates ranging between r_1 and r_2 and thus

$$\mathbb{E}_{(X_{t_m}, K_{t_m})} \left[\left| \langle M \rangle_{t_{m+1}} - \langle M \rangle_{t_m} - \sigma(t_{m+1} - t_m) \right| \right] \leq |r_2 - \sigma| \vee |r_1 - \sigma| \mathcal{R}^{-1} < r_2 \mathcal{R}^{-1}. \quad (7.4)$$

To bound $t^{-1} \mathcal{R}^{-1} r_2 \mathbb{E}[(1 - \chi(S_{\epsilon,\delta}^{(t)})) \mathcal{N}_t]$, we apply the Cauchy-Schwarz inequality to attain $2^{\frac{1}{2}} t \mathcal{R} (1 - \Pr[S_{\epsilon,\delta}^{(t)}])^{\frac{1}{2}}$.

In the event $S_{\epsilon,\delta}^{(t)}$, by the discussion in the second paragraph of this proof, we expect that a fraction of at least $(1 - \delta)$ of the times t_n have $|K_{t_n}^-| \geq \epsilon t^{\frac{1}{2}}$. For large t , it will be easy to see that the terms corresponding to momentum jumps $K_{t_n} - K_{t_n}^- = v_n$ with $|v_n| \geq \frac{1}{2} \epsilon t^{\frac{1}{2}}$ will be negligible, and we can assume $|K_{t_n}| > \frac{1}{2} \epsilon t^{\frac{1}{2}}$. Using that the differences $t_{n+1} - t_n$ are exponentially distributed with expectation $\mathcal{R}^{-1}$, then for $(x, k) = (X_{t_n}, K_{t_n})$, $|k| \geq 2^{-1} \epsilon t^{\frac{1}{2}}$

$$\begin{aligned} \mathbb{E}_{(x,k)} \left[\left| \langle M \rangle_{t_{n+1}} - \langle M \rangle_{t_n} - \sigma(t_{n+1} - t_n) \right| \right] &= \int_0^\infty d\tau \mathcal{R} e^{-\mathcal{R}\tau} \left| \int_0^1 da (T_{(x,k),\tau}(a) - \tau) \int_{\mathbb{R}} dv j_a(v) v^2 \right| \\ &\leq r_2 |k|^{-1} \int_0^\infty d\tau \mathcal{R} \tau e^{-\mathcal{R}\tau} \leq 2 r_2 \mathcal{R}^{-1} \epsilon^{-1} t^{-\frac{1}{2}}, \end{aligned} \quad (7.5)$$

where $r_2 = \sup_{0 \leq a \leq 1} \int_{\mathbb{R}} dv j_a(v) v^2$ and $T_{(x,k),\tau} \in L^1([0, 1])$ is as defined in Lemma 5.2 which has been applied to get the second inequality.

Our recipe for bounding (7.2) is the following:

1. Pick δ so that $r_2 \delta \ll 1$,

2. Pick ϵ so that $r_2(C\frac{\sqrt{r_2}\epsilon}{r_1\delta})^{\frac{1}{2}} \ll 1$,
3. Pick t so that $r_2\epsilon^{-1}t^{-\frac{1}{2}} \ll 1$.

By the observations in the second paragraph of this proof, $1 - \Pr[S_{\epsilon,\delta}^{(t)}] \leq C\frac{\sqrt{r_2}\epsilon}{r_1\delta}$, and hence the first term on the right side of (7.2) is small.

Let n'_ϵ be the number of Poisson times with $|K_{t_n^-}| > 2^{-1}\epsilon t^{\frac{1}{2}}$ and $|K_{t_n}| \leq 2^{-1}\epsilon t^{\frac{1}{2}}$ up to time t , and n''_ϵ be the number of times with $|K_{t_n}| \geq 2^{-1}\epsilon t^{\frac{1}{2}}$.

Applying the bounds (7.5) and (7.4) depending on whether or not $|K_{t_n}| \geq 2^{-1}\epsilon t^{\frac{1}{2}}$, then we get the first inequality below:

$$\begin{aligned} \mathbb{E}_\mu \left[\chi(S_{\epsilon,\delta}^{(t)}) \frac{1}{t} \sum_{m=0}^{\mathcal{N}_t} \mathbb{E}_{(X_{t_m}, K_{t_m})} [|\langle M \rangle_{t_{m+1}} - \langle M \rangle_{t_m} - \sigma(t_{m+1} - t_m)|] \right] \\ < \mathbb{E}_\mu \left[\chi(S_{\epsilon,\delta}^{(t)}) ((n_{\epsilon,t} + n'_\epsilon) \mathcal{R}^{-1} r_2 t^{-1} + n''_\epsilon 2r_2 \mathcal{R}^{-1} \epsilon^{-1} t^{-\frac{3}{2}}) \right] \leq r_2 \delta + O(t^{-\frac{1}{2}}) \ll 1. \end{aligned} \quad (7.6)$$

Now we look at the expectation of the terms $n_{\epsilon,t}$, n'_ϵ , and n''_ϵ to reach the second inequality. Since n''_ϵ is smaller than the total number of Poisson times $\mathcal{N}_t$, $\mathbb{E}_\mu[n''_\epsilon] \leq \mathcal{R}t$, which makes its contribution in (7.6) $O(t^{-\frac{1}{2}})$. The number $n'_{\epsilon,t}$ is smaller than $\sum_{n=1}^{\mathcal{N}_t} \chi(|v_n| \geq 2^{-1}\epsilon t^{\frac{1}{2}})$, so for the distribution $\mathcal{P}_a(v)$ of a momentum jump conditioned to occur at a torus point $a \in [0, 1]$, we can write

$$t^{-1} \mathbb{E}_\mu[n'_{\epsilon,t}] \leq t^{-1} \mathbb{E}_\mu \left[\sum_{n=1}^{\mathcal{N}_t} \chi(|v_n| \geq 2^{-1}\epsilon t^{\frac{1}{2}}) \right] \leq 2t^{-1} \mathbb{E}_\mu[\mathcal{N}_t] \sup_{0 \leq a \leq 1} \int_{2^{-1}\epsilon t^{\frac{1}{2}}}^{\infty} dv \mathcal{P}_a(v) = O(t^{-1}).$$

The decay for the above expression follows by Chebyshev's inequality and by the uniformly bounded second moments for $\mathcal{P}_a$, $a \in [0, 1]$. To bound the $t^{-1}n_{\epsilon,t}$, we will finally be able to use the definition of the event $S_{\epsilon,\delta}$ in the second inequality below

$$\mathbb{E}_\mu [\chi(S_{\epsilon,\delta}^{(t)}) t^{-1} n_{\epsilon,t}] \leq \mathcal{R} \mathbb{E}_\mu [\chi(S_{\epsilon,\delta}^{(t)}) \int_0^1 ds \chi(t^{-\frac{1}{2}} |K_{st}| \leq \epsilon)] + O(t^{-\frac{1}{2}}) \leq \mathcal{R} \delta + O(t^{-\frac{1}{2}}).$$

Finally, the first inequality follows from (7.1). □

Proof of Theorem 4.1. By Lemma 4.2, $t^{-\frac{1}{2}}|K_{st}|$ obeys the equation

$$\begin{aligned} t^{-\frac{1}{2}}|K_{st}| &= \int_0^s S(K_{rt}) dM_r^{(t)} + \sup_{0 \leq a \leq s} - \int_0^a S(K_{rt}) dM_r^{(t)} + \mathcal{E}_s^{(t)} \\ &= t^{-\frac{1}{2}} \sum_{n=1}^{\mathcal{N}_{st}} w_n S(K_{t_n^-}) + t^{-\frac{1}{2}} \sup_{0 \leq s \leq 1} - \sum_{n=1}^{\mathcal{N}_{st}} w_n S(K_{t_n^-}) + \mathcal{E}_s^{(t)} \end{aligned} \quad (7.7)$$

where the error $\mathcal{E}_s^{(t)}$ vanishes in the norm $\mathbb{E}[\sup_{0 \leq s \leq 1} |\mathcal{E}_s^{(t)}|]$. The martingale $M'_r = \sum_{n=1}^{\mathcal{N}_r} w_n S(K_{t_n^-})$ has the same quadratic variation as the martingale of momentum jumps M_r .

$$[M]_r = [M']_r = \sum_{n=1}^{\mathcal{N}_r} w_n^2.$$

Thus the predictable quadratic variations $\langle M \rangle_r$ and $\langle M' \rangle$ are also equal.

However, by the proof of Theorem 3.1, $\mathbb{E}_\mu[|t^{-1}\langle M \rangle_{st} - \sigma s|] \rightarrow 0$ for $s \in [0, 1]$, and thus M'_r converges to a Brownian motion. By the same argument as at the end of the proof of Lemma 3.2, we have that $t^{-\frac{1}{2}}|K_{st}|$ converges to the absolute value of a Brownian motion. $\square$

Proof of Main Result. It is sufficient to prove that $t^{-\frac{1}{2}}K_{st}$ converges to a Brownian motion, since by a change of integration variable

$$t^{-\frac{3}{2}}X_{st} = t^{-\frac{3}{2}}\left(X_0 + \int_0^{st} dr K_r\right) = t^{-\frac{3}{2}}X_0 + \int_0^s dr t^{-\frac{1}{2}}K_{rt}.$$

Hence $(t^{-\frac{3}{2}}X_{st}, t^{-\frac{1}{2}}K_{st})$ converges in distribution to $(\int_0^s dr \mathbf{B}_r, \mathbf{B}_s)$ if $t^{-\frac{1}{2}}K_{st}$ converges to $\mathbf{B}_s$.

Recall that $K_t = K_0 + M_t + \int_0^t ds \frac{dV}{dx}(X_s)$. By Lemma 6.8, the first moment of

$$\sup_{0 \leq s \leq t} \left| t^{-\frac{1}{2}} \int_0^s dr \frac{dV}{dx}(X_r) \right|$$

converges to zero. By Theorem 3.1, $t^{-\frac{1}{2}}M_{st}$ converges to a Brownian motion with diffusion constant σ . Hence $t^{-\frac{1}{2}}K_{st}$ converges to a Brownian motion. $\square$

Acknowledgments

We thank Wojciech De Roeck, Karel Netočný and Frank Redig for useful discussions. J. C. acknowledges support from the Belgian Interuniversity Attraction Pole P6/02 and the Marie Curie Research Training Network project MRTN-CT-2006-035651, Acronym CODY, of the European Commission.

A A martingale lemma

Below we formalize a simple lemma for a martingale $\mathbf{M}_t$ which makes jumps at discrete times. For a basic introduction to martingale theory, see [3].

Lemma A.1. *Let $\mathbf{M}_t = \sum_{n=1}^{\mathbf{N}_t} X_n$ be a right-continuous martingale adapted to a filtration $\mathcal{F}_t$, and making jumps $X_n(\omega)$, $\omega \in \mathcal{F}_{t_n}$ at discrete times according to some random counter $\mathbf{N}_t$ where $t_n = \inf\{s \in \mathbb{R}^+ | \mathbf{N}_s = n\}$. Assume also that $\mathbb{E}[\mathbf{N}_t] < \infty$ for every t and that $\mathbb{E}[X_n^2] < \infty$ for every n . Then*

$$\mathbb{E}[\mathbf{M}_t^2] \leq \mathbb{E}[\mathbf{N}_t] \sup_n \mathbb{E}[X_n^2 | n \leq \mathbf{N}_t]$$

where $\mathbb{E}[X_n^2 | n \leq \mathbf{N}_t]$ is defined as zero for n such that $\Pr[n \leq \mathbf{N}(t)] = 0$.

Proof. By orthogonality of martingale increments

$$\begin{aligned} \mathbb{E}[\mathbf{M}_t^2] &= \mathbb{E}\left[\sum_{n=1}^{\mathbf{N}_t} X_n^2\right] = \mathbb{E}\left[\sum_{n=1}^{\infty} X_n^2 \chi(n \leq \mathbf{N}_t)\right] = \sum_{n=1}^{\infty} \Pr[n \leq \mathbf{N}_t] \mathbb{E}[X_n^2 | n \leq \mathbf{N}_t] \\ &\leq \left(\sum_{n=1}^{\infty} \Pr[n \leq \mathbf{N}_t]\right) \sup_n \mathbb{E}[X_n^2 | n \leq \mathbf{N}_t] \leq \mathbb{E}[\mathbf{N}_t] \sup_n \mathbb{E}[X_n^2 | n \leq \mathbf{N}_t], \end{aligned}$$

where the inequality is Hölder's. $\square$

B Boundary crossing distributions

Consider symmetric independent and identically distributed random variables $X_1, X_2, \dots$ with mean zero and density

$$\tilde{P}(v) = \int_0^1 da \frac{\kappa(a)}{\bar{\kappa}} \mathcal{P}_a(v).$$

Construct the random walk $Y_n = \sum_{m=1}^n X_m$. We will refer to this as the so called “averaged random walk” Let $L \geq 0$. Our interest in this section is to understand the probability density

$$\pi_L(a, b) = \sum_{n=1}^{\infty} \Pr[Y_n - L = a, X_n = b, Y_r < L \text{ for } r < n], \quad a, b \leq \mathbb{R}^+. \quad (\text{B.1})$$

In the present appendix we use the notation $\Pr[\cdot]$ for the induced density by the random walk, in its obvious meaning for continuous densities. To be clear, π_L describes the distribution of jumps in excess over point L for the random walk Y_n on the first time that it passes over that L .

For $L = 0$, we write $D(v, w) = \pi_{L=0}(v, w)$ which is in fact the “successive record increment” distribution. Indeed, define $R_n = \sup_{0 \leq m \leq n} Y_m$, the record in the positive direction for the random walk up to time n , and let τ_m be the time of the m th record. Then, the increments $R_{\tau_m} - R_{\tau_{m-1}}$ are i.i.d. random variables with density $D(a) = \int_0^\infty db D(a, b)$. We use this fact in the proof of the following proposition. In particular

$$\mathbb{E}[R_{\tau_m} - R_{\tau_{m-1}}] = \int_0^\infty \int_0^\infty da db a D(a, b).$$

The topic of “record distributions” has a much wider scope and has a long history in extreme value statistics, see e.g. [8] for some pioneering contribution.

Lemma B.1. *Assume (III) of List 2.1 and (i)-(ii) of List 2.2.*

1. *The Laplace transform $\varphi(q)$ of $D(v)$ satisfies*

$$\varphi(q) \leq \frac{\mathcal{C}}{1 - e^{-\eta - q}} \quad \text{for } q > -\eta.$$

2. *$D(v)$ is bounded and continuous.*

3. *The following is a probability density on $\mathbb{R}^+ \times \mathbb{R}^+$*

$$\pi_\infty(v, w) = \frac{\int_v^\infty dx D(x, w)}{\int_0^\infty \int_0^\infty dx dy x D(x, y)},$$

which is positive for all $v < w$.

Proof. Part (1) follows, since $D(v)$ inherits the property (i) of List 2.2. □

Proposition B.2. *Assume (III) of List 2.1 and (i)-(ii) of List 2.2. Let the random walk Y_n , $\pi_L(v, w)$, and $D(v, w)$ be defined as above.*

In the limit of large L , we have the $L^1(\mathbb{R}^+ \times \mathbb{R}^+)$ convergence

$$\pi_L(v, w) \longrightarrow \pi_\infty(v, w) = \frac{\int_v^\infty dx D(x, w)}{\int_0^\infty \int_0^\infty dx dy x D(x, y)}$$

Moreover, the difference between π_∞ and π_L converges exponentially and monotonically.

Corollary B.3. *Let Y_n be a random walk as above. For $L > 0$ and $d \in \mathbb{R}$ with $|d| \leq \frac{1}{2}L$ define $S = (-d - L, L) \subset \mathbb{R}$, and let $\pi_{\uparrow, L}(v, w), \pi_{\downarrow, L}(v, w)$ be the probability densities on $\mathbb{R}^+ \times \mathbb{R}^+$ defined as*

$$\pi_{\uparrow, L}(v, w) = \sum_{n=1}^{\infty} \Pr[X_n = w, Y_n - L = v, Y_m \in S \text{ for } 0 \leq m < n] 1_{Y_n > 0}. \quad (\text{B.2})$$

$$\pi_{\downarrow, L}(v, w) = \sum_{n=1}^{\infty} \Pr[-X_n = w, Y_n + L + d = -v, Y_m \in S \text{ for } m < n] 1_{Y_n < 0} \quad (\text{B.3})$$

In the limit $L \rightarrow \infty$, we have $L^1(\mathbb{R}^+ \times \mathbb{R}^+)$ convergence

$$\pi_{\uparrow, t} \longrightarrow p_{\uparrow} \pi_{\infty} \quad \pi_{\downarrow, t} \longrightarrow p_{\downarrow} \pi_{\infty}$$

Moreover, the convergence is uniform for $|d| \leq \frac{1}{2}L$.

Proof. In general, we have the identity

$$\pi_L(v, w) = \pi_{\uparrow, L}(v, w) + \int_{\mathbb{R}^+ \times \mathbb{R}^+} da db \pi_{\downarrow}(a, b) \pi_{2L+d+b}(v, w), \quad (\text{B.4})$$

where $\pi_L(v, w)$ is the boundary increment distribution from (B.2). As $L \rightarrow \infty$, $\pi_L(v, w), \pi_{2L+d+b}(v, w)$ converge exponentially in $L^1(\mathbb{R}^+ \times \mathbb{R}^+)$ to $\pi_{\infty}(v, w)$ by Proposition B.2. By the triangle inequality and rearranging (B.4),

$$\|\pi_{\uparrow, L} - \pi_{\infty}\|_1 \leq \int_{\mathbb{R}^+ \times \mathbb{R}^+} da db \pi_{\downarrow}(a, b) \|\pi_{2L+d+b} - \pi_L\|_1 \leq 2 p_{\downarrow} \sup_{L' \geq L} \|\pi_{L'} - \pi_{\infty}\|_1,$$

where $p_{\uparrow} = \int_{\mathbb{R}^+ \times \mathbb{R}^+} dv dw \pi_{\uparrow, L}(v, w)$ and $p_{\downarrow} = \int_{\mathbb{R}^+ \times \mathbb{R}^+} dv dw \pi_{\downarrow, L}(v, w)$. Since $|d| \leq \frac{1}{2}L$, $p_{\uparrow}^{-1}$ is bounded away from zero and the convergence holds.

The uniformity of the convergence over $|d| \leq \frac{1}{2}L$ is implied by the monotonicity of the convergence from Proposition B.2. □

Next we extend Proposition B.2 and Corollary B.3 to results about our dynamics in high momentum situations. We prove an analogue of Corollary B.3 and then prove the analogue of Proposition B.2 as a corollary. Since there is also a position variable moving on the torus, the crossing distributions are joint distributions on $[0, 1] \times \mathbb{R}^+$. We consider the so called “averaged random walk”

$$\tilde{P}(v) = \int_0^1 da \frac{\kappa(a)}{\bar{\kappa}} \mathcal{P}_a(v),$$

where $\bar{\kappa} = \int_0^1 da \kappa(a)$, and its boundary crossing distribution $\pi_{\infty}(v, w)$ obtained from the previous Proposition B1 gives rise to another density

$$\phi_{\infty}(a, v) = \int_{\mathbb{R}^+} dw \pi_{\infty}(v, w) \frac{\kappa(a) \mathcal{P}_a(w)}{\int_0^1 dx \kappa(x) \mathcal{P}_x(w)}. \quad (\text{B.5})$$

Intuitively, $\phi_{\infty}(a, v)$ is an idealized joint distribution for the position on the torus and the over-jump for the particle upon crossing the border of some domain $S \subset [0, 1] \times \mathbb{R}^+$. $\phi_{\infty}(a, v)$ provides a good approximation for such a distribution at high energy where the momentum behaves nearly as the averaged random walk.

The following lemma is similar to Part (1) of Lemma (B.1).

Lemma B.4. Assume (i) of List 2.2. Let $S \subset \mathbb{R}$ be an open interval with boundaries $L_1 < L_2$ not both unbounded. If the particle begins at some point (x, k) with $k \in S$, then define

$$\begin{aligned}\pi_{\uparrow}(v) &= \sum_{n=1}^{\infty} \Pr[K_{t_n} - L_2 = v \geq 0, K_{t_m} \in S \text{ for } 0 \leq m < n] 1_{K_{t_n} \geq S}, \\ \pi_{\downarrow}(v) &= \sum_{n=1}^{\infty} \Pr[K_{t_n} - L_1 = -v \leq 0, K_{t_m} \in S \text{ for } 0 \leq m < n] 1_{K_{t_n} \leq S}.\end{aligned}$$

There exists a universal upper bound J such that for all L_1, L_2 and (x, k)

$$\frac{\int_{\mathbb{R}^+} dv \pi_{\uparrow}(v) v^2}{\int_{\mathbb{R}^+} dv \pi_{\uparrow}(v)}, \frac{\int_{\mathbb{R}^+} dv \pi_{\downarrow}(v) v^2}{\int_{\mathbb{R}^+} dv \pi_{\downarrow}(v)} \leq J,$$

when $\int_{\mathbb{R}^+} dv \pi_{\uparrow}(v), \int_{\mathbb{R}^+} dv \pi_{\downarrow}(v) \neq 0$. More generally, the Laplace transforms $\varphi_{\uparrow}, \varphi_{\downarrow}$ have

$$\varphi_{\uparrow}(q), \varphi_{\downarrow}(q) \leq \frac{\mathcal{C}}{1 - e^{-\eta - q}} \quad \text{for } q > -\eta.$$

Let $P(v_1, \dots, v_n)$ be the joint probability distribution for the increments $v_j = K_{t_j} - K_{t_{j-1}}$ where t_j is the time of the j th momentum jump.

Lemma B.5. Let us make assumptions (i)-(ii) from List 2.2. Fix $\beta, \gamma > 0$. Let the dynamics begin at a point (x, k) and $v_1, \dots, v_n \in [-2t^\gamma, 2t^\gamma]$ be such that $|k + \sum_{r=0}^m v_r| \geq \frac{1}{2}t^\beta$ for all $0 \leq m \leq n$. Then

$$\left| \frac{P(v_1, \dots, v_n)}{\tilde{P}(v_1) \dots \tilde{P}(v_n)} - 1 \right| < c n t^{\gamma - \beta} e^{c n t^{\gamma - \beta}},$$

where $c = 8\bar{V}\mu$.

Proof. We start with the one-jump case. The only difference between Lemma B.5 and Lemma 5.2 is that here the increment v is the sum of the first momentum kick and the drift up to the time of the kick.

Let (x_s, k_s) be on the trajectory determined by the Hamiltonian $H(x, k) = \frac{1}{2}k^2 + V(x)$ starting from $(x, k) \in [0, 1] \times \mathbb{R}$. For fixed starting point (x, k) , $\Delta_s = k_s - k = \Delta(x_s)$ is function of x_s . For the one-jump case, we can use the density $\tilde{r}_{(x,k)} \in L^1([0, 1])$ from the proof of Lemma 5.2, to write

$$P(v) = \int_0^1 da \tilde{r}_{(x,k)}(a) \mathcal{P}_a(v - \Delta(a))$$

Define the density $P'(v) = \int_0^1 da \frac{\kappa(a)}{\bar{\kappa}} \mathcal{P}_a(v - \Delta(a))$. Following the same line of reasoning as (5.6) in Lemma 5.2,

$$\begin{aligned}\sup_{v \in \mathbb{R}} \left| \frac{P(v)}{P'(v)} - 1 \right| &\leq \sup_{v \in \mathbb{R}} \frac{1}{P'(v)} \int_0^1 da \left| \tilde{r}_{(x,k)}(a) - \frac{\kappa(a)}{\bar{\kappa}} \right| \mathcal{P}_a(v - \Delta(a)) \\ &\leq \sup_{0 \leq a \leq 1} \left| \tilde{r}_{(x,k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \frac{1}{P'(v)} \int_0^1 da \frac{\kappa(a)}{\bar{\kappa}} \mathcal{P}_a(v - \Delta(a)) = \sup_{a \in [0, 1]} \left| \tilde{r}_{(x,k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right|\end{aligned}$$

Since, $|k| \geq \frac{1}{2}t^\beta$, then by Part (2) of Lemma 5.2,

$$\sup_{a \in [0, 1]} \left| \tilde{r}_{(x,k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \leq \nu^{-2} 4\mathcal{R} t^{-\beta} + O(t^{-2\beta}).$$

Moreover,

$$\begin{aligned}
\left| \frac{P'(v)}{\tilde{P}(v)} - 1 \right| &\leq \frac{1}{\tilde{P}(v)} \int_0^1 da \frac{\kappa(a)}{\bar{\kappa}} |\mathcal{P}_a(v - \Delta(a)) - \mathcal{P}_a(v)| \\
&= \frac{1}{\tilde{P}(v)} \int_0^1 da \frac{\kappa(a)}{\bar{\kappa}} |\Delta(a)| \left| \int_0^1 ds \frac{d\mathcal{P}_a}{dv}(v + s\Delta(a)) \right| \\
&\leq \frac{\mu 4\bar{V}(1 + |v|)t^{-\beta}}{\tilde{P}(v)} \int_0^1 da \frac{\kappa(a)}{\bar{\kappa}} \mathcal{P}_a(v) = \mu 4\bar{V}(1 + |v|)t^{-\beta} \quad (\text{B.6})
\end{aligned}$$

For the second inequality we have used Lemma 5.1 to get $\sup_{0 \leq s \leq 1} |\Delta(a)| \leq 4\bar{V}t^{-\beta}$. In particular, $|\Delta(a)| < 1$ allows us to use (ii) of List 2.2 to bound $|\frac{d\mathcal{P}_a}{dv}(v + s\Delta(a))|$ by $\mu(1 + |v|)\mathcal{P}_a(v)$. Since $1 + |v| \leq 1 + 2t^\gamma < 4t^\gamma$ for large enough t , we have

$$\sup_{v \in \mathbb{R}} \left| \frac{P'(v)}{\tilde{P}(v)} - 1 \right| < 16\bar{V}\mu t^{\gamma-\beta}$$

$$\begin{aligned}
\left| \frac{P(v)}{\tilde{P}(v)} - 1 \right| &\leq \left| \frac{P(v)}{P'(v)} - 1 \right| \left| \frac{P'(v)}{\tilde{P}(v)} \right| + \left| \frac{P'(v)}{\tilde{P}(v)} - 1 \right| \leq (16\bar{V}\mu t^{\gamma-\beta}) (1 + 4\mathcal{R}\nu^{-2}t^{-\beta} + O(t^{-2\beta})) \\
&\quad + 4\mathcal{R}\nu^{-2}t^{-\beta} + O(t^{-2\beta}) < 20\bar{V}\mu t^{\gamma-\beta} = ct^{\gamma-\beta} \quad (\text{B.7})
\end{aligned}$$

where the last inequality is for t large enough and $c = 20\bar{V}\mu$.

To extend to arbitrary n

$$\left| \frac{P(v_1, \dots, v_n)}{\tilde{P}(v_1) \dots \tilde{P}(v_n)} - 1 \right| \leq \sum_{m=1}^n \left| \frac{P(v_1, \dots, v_m)}{\tilde{P}(v_1) \dots \tilde{P}(v_m)} - \frac{P(v_1, \dots, v_{m-1})}{\tilde{P}(v_1) \dots \tilde{P}(v_{m-1})} \right| \quad (\text{B.8})$$

Moreover, we can write the summand as

$$\left| \frac{P(v_1, \dots, v_m)}{P(v_1, \dots, v_{m-1}) \tilde{P}(v_m)} - 1 \right| \prod_{r=1}^{m-1} \frac{P(v_1, \dots, v_r)}{P(v_1, \dots, v_{r-1}) \tilde{P}(v_r)}$$

The ratio's appearing in the above expression can be written in terms of an expectation:

$$\frac{P(v_1, \dots, v_m)}{P(v_1, \dots, v_{m-1}) \tilde{P}(v_m)} = \mathbb{E} \left[\frac{P_{(X_{t_{m-1}}, K_{t_{m-1}})}(v_m)}{\tilde{P}(v_m)} \middle| K_{t_r} - K_{t_{r-1}} = v_r \text{ for } r < m \right] \quad (\text{B.9})$$

where $P_{(X_{t_{m-1}}, K_{t_{m-1}})}(v_m)$ is the one jump distribution starting from $(x, k) = (X_{t_{m-1}}, K_{t_{m-1}})$. The right side of (B.9) is thus an average of the one-jump case over different starting points $(x, k) = (X_{t_{m-1}}, K_{t_{m-1}})$ and thus we can apply the $n = 1$ case.

We thus conclude

$$\left| \frac{P(v_1, \dots, v_n)}{\tilde{P}(v_1) \dots \tilde{P}(v_n)} - 1 \right| < cn t^{\gamma-\beta} (1 + ct^{\gamma-\beta})^n < cn t^{\gamma-\beta} e^{cn t^{\gamma-\beta}}.$$

□

For notational convenience, we will take the initial point (x, k) to have positive momentum $k > 0$ for the propositions and lemmas below.

Proposition B.6. Assume (i)-(ii) from List 2.2. Let (X_s, K_s) evolve from an initial point (x, k) with $k \geq t^\beta$. Define $S = \{y \in \mathbb{R} \mid k - t^\gamma - d < y < k + t^\gamma\}$ for $0 < \gamma < \frac{1}{4}\beta$ and $|d| \leq \frac{1}{2}t^\gamma$, and the densities

$$\begin{aligned}\pi_{\uparrow, t}(v, w) &= \sum_{n=1}^{\infty} \Pr[K_{t_n} - K_{t_{n-1}} = w, K_{t_n} - k - t^\gamma = v \geq 0, K_{t_m} \in S \forall (m < n)] 1_{K_{t_n} > S} \\ \pi_{\downarrow, t}(v, w) &= \sum_{n=1}^{\infty} \Pr[K_{t_n} - K_{t_{n-1}} = w, K_{t_n} - k + t^\gamma + d = -v \leq 0, K_{t_m} \in S \forall (m < n)] 1_{K_{t_n} < S}\end{aligned}$$

In the limit $t \rightarrow \infty$, there is $L^1(\mathbb{R}^+ \times \mathbb{R}^+)$ convergence

$$\pi_{\uparrow, t} \longrightarrow p_{\uparrow} \pi_{\infty}, \quad \pi_{\downarrow, t} \longrightarrow p_{\downarrow} \pi_{\infty},$$

where $p_{\uparrow}$ and $p_{\downarrow}$ are the exit probabilities for the averaged random walk, and the convergence is uniform for all $|d| \leq \frac{1}{2}t^\gamma$.

Proof. Define the sets

$$S_n(p, w) = \{(v_1, \dots, v_n) \in \mathbb{R}^n \mid v_n = w, p = k + \sum_{r=1}^n v_r \notin S, k + \sum_{r=1}^m v_r \in S \text{ for } m < n\}.$$

and define $S'_n(p, w)$ so that $S'_n(p, w) = S_n(p, w) \cap \{|v_m| \leq 2t^\gamma \text{ for } m \leq n\}$ when $n \leq \lfloor t^{3\gamma} \rfloor$ and $S'_n(p, w) = \emptyset$ when $n > \lfloor t^{3\gamma} \rfloor$.

Then $\pi_{\uparrow, t}(v, w), \pi_{\downarrow, t}(v, w)$ satisfy

$$\pi_{\uparrow, t}(p - k - t^\gamma, w) + \pi_{\downarrow, t}(k - t^\gamma - d - p, w) = \sum_{n=1}^{\infty} \int_{S_n(p, w)} dv_1 \cdots dv_n P(v_1, \dots, v_n) \quad (\text{B.10})$$

where $P(v_1, \dots, v_n)$ are the joint densities from Lemma B.5. The sum of $\pi_{\uparrow, t}$ and $\pi_{\downarrow, t}$ forms a probability measure, since, by an analogous argument as in the proof of Lemma 6.1, the exit time for S is almost surely finite (and has expectation $\propto t^{2\beta}$). We will argue that the right side is close in $L^1(\mathbb{R}^+ \times \mathbb{R}^+)$ as a function of (p, w) to the same expression with $P(v_1, \dots, v_n)$ replaced by $\tilde{P}(v_1) \cdots \tilde{P}(v_n)$:

$$\pi_{\uparrow, t}^{(0)}(p - k - t^\gamma, w) + \pi_{\downarrow, t}^{(0)}(k - t^\gamma - d - p, w) = \sum_{n=1}^{\infty} \int_{S_n(p, w)} dv_1 \cdots dv_n \tilde{P}(v_1) \cdots \tilde{P}(v_n).$$

This expression corresponds to the averaged random walk. We can then apply Lemma B.3 for an ordinary random walk to get the convergence of $\pi_{\uparrow, t}^{(0)}$ and $\pi_{\downarrow, t}^{(0)}$ to $p_{\uparrow} \pi_{\infty}$ and $p_{\downarrow} \pi_{\infty}$.

By definition of $S_n(p, w)$, $S'_n(p, w)$, $K + \sum_{m=1}^n v_m > \frac{1}{2}t^\beta$ for all $0 \leq m < n$. First, notice that by Lemma B.5

$$\begin{aligned}\sum_{n=1}^{\infty} \int_{S'_n(p, w)} dv_1 \cdots dv_n \tilde{P}(v_1) \cdots \tilde{P}(v_n) \left| \frac{P(v_1, \dots, v_n)}{\tilde{P}(v_1) \cdots \tilde{P}(v_n)} - 1 \right| \\ \leq \sum_{n=1}^{\lfloor t^{3\gamma} \rfloor} \int_{S'_n(p, w)} dv_1 \cdots dv_n \tilde{P}(v_1) \cdots \tilde{P}(v_n) c n t^{\gamma-\beta} e^{c n t^{\gamma-\beta}} \quad (\text{B.11})\end{aligned}$$

The $L^1(\mathbb{R}^+ \times \mathbb{R}^+)$ norm of the right-side of (B.11) (by integrating over p, w) is bounded by $\mathbb{E}[q(N_0 \wedge t^{3\gamma}) e^{q(N_0 \wedge t^{3\gamma})}]|_{q=ct^{\gamma-\beta}}$, where N_0 is the number of steps that a random walk with jumps having density $c_t^{-1} \tilde{P}(v) 1_{|v| \leq t^\gamma}$, for normalization constant c_t , takes to leave S starting from k . The right-side of (B.11) vanishes as

$$\mathbb{E}[q(N_0 \wedge t^{3\gamma}) e^{q(N_0 \wedge t^{3\gamma})}]|_{q=ct^{\gamma-\beta}} \leq c t^{4\gamma-\beta} e^{c t^{4\gamma-\beta}} \approx c t^{4\gamma-\beta}.$$

Now we need to show that not much probability was lost by replacing $S_n(p, w)$ with $S'_n(p, w)$. Since both $\pi_{\uparrow, t} + \pi_{\downarrow, t}$ and $\pi_{\uparrow, t}^{(0)} + \pi_{\downarrow, t}^{(0)}$ are probability measures, it is enough to show that the probability of the event $\{|v_m| > 2t^\gamma \text{ for some } m \leq N \text{ or } N > t^{3\gamma}\}$ is small for the random walk. First, note that the expected number of steps N to leave S for the averaged random walk will be smaller than the expected number of the steps N_0 for the capped random walk.

$$\Pr[N > t^{3\gamma}] \leq \Pr[N_0 > t^{3\gamma}] \text{ and } \mathbb{E}[N] \leq \mathbb{E}[N_0].$$

This follows since the first jump of size greater than $t^{2\gamma}$ will immediately leave S , and the corresponding capped trajectory may have to continue on for more steps before leaving S .

Now we argue that $\mathbb{E}[N_0] = O(t^{2\gamma})$. Let us set $k = 0$. Define the stopping time $N_{0, \mathcal{T}} = N_0 \wedge \mathcal{T}$. Reasoning as in the proof of Lemma 6.1, then

$$\mathbb{E}[N_{0, \mathcal{T}}] = \zeta^{-1} \mathbb{E}\left[\sum_{n=1}^{N_{0, \mathcal{T}}} v_n^2\right] = \zeta^{-1} \mathbb{E}[(v_1 + \dots + v_{N_{0, \mathcal{T}}})^2] \leq \zeta^{-1} (3t^\gamma + d)^2 < 16 \zeta^{-2} t^{2\gamma},$$

where the expectations are with respect to the statistics for the capped random walk, and $\xi = c_t^{-1} \int_{|v| \leq t^\gamma} dv \tilde{P}(v) v^2$.

In the second equality, we used that $v_1 + \dots + v_{N_{0, \mathcal{T}}}$ is either inside $[-t^\gamma - d, t^\gamma]$ when $\mathcal{T} < N_0$ or has jumped out this interval with a jump smaller than $t^{2\gamma}$. The bound on the right is independent of $\mathcal{T}$, so

$$\mathbb{E}[N_0] = \limsup_{\mathcal{T} \rightarrow \infty} \mathbb{E}[N_{0, \mathcal{T}}] \leq 16 \zeta^{-2} t^{2\gamma}.$$

By Chebyshev's inequality

$$\Pr[N > t^{3\gamma}] \leq t^{-3\gamma} \mathbb{E}[N_0] \leq 16 \zeta^{-1} t^{-\gamma} \rightarrow 0.$$

We still need to show that $\Pr[|v_n| > 2t^\gamma \text{ for some } n \leq N]$ is small.

$$\Pr[|v_n| > 2t^\gamma \text{ for some } n \leq N] \leq \mathbb{E}[N] \Pr[|v_n| \geq t^\gamma].$$

By our remarks above $\mathbb{E}[N] \leq \mathbb{E}[N_0] = O(t^{2\gamma})$. Using Chebyshev's inequality, and the bound on the fourth moment of a single momentum jump by ρ ,

$$t^{4\gamma} \Pr[|v_n| \geq t^{2\gamma}] \leq \mathbb{E}[v_n^4] \leq \rho.$$

Putting the above inequalities together

$$\Pr[|v_n| \geq t^\gamma, \text{ for some } n \leq N] \leq \mathbb{E}[N] \Pr[|v_n| \geq t^\gamma] < 16 \rho \zeta^{-1} t^{-2\gamma}.$$

Hence the event that the last jump v_n

We have shown that $\pi_{\uparrow, t}, \pi_{\downarrow, t}$ converge to π_∞ in $L^1(\mathbb{R}^+ \times \mathbb{R}^+)$ at $t \rightarrow \infty$. The convergence of $\pi_{\uparrow, t}$ and $\pi_{\downarrow, t}$ to $\pi_{\infty, t}$ is uniform over $|d| \leq \frac{1}{2}t^\gamma$ by the uniformity in Lemma B.3 and by the uniformity in the bounds above.

□

Proposition B.7. Assume (i)-(ii) from List 2.2. Let (X_s, K_s) evolve from an initial point (x, k) with $k \geq t^\beta$. Define $S = \{y \in \mathbb{R} \mid k - t^\gamma - d < y < k + t^\gamma\}$ for $0 < \gamma < \frac{1}{4}\beta$, and the densities

$$\begin{aligned}\phi_{\uparrow, t}(a, v) &= \sum_{n=1}^{\infty} \Pr[X_{t_n} = a, K_{t_n} - k - t^\gamma = v \geq 0, K_{t_m} \in S \text{ for } 0 \leq m < n] 1_{K_{t_n} > S}, \\ \phi_{\downarrow, t}(a, v) &= \sum_{n=1}^{\infty} \Pr[X_{t_n} = a, K_{t_n} - k + t^\gamma + d = -v \leq 0, \text{ for } K_{t_m} \in S \text{ } 0 \leq m < n] 1_{K_{t_n} < S}.\end{aligned}$$

In the limit $t \rightarrow \infty$, there is $L^1([0, 1] \times \mathbb{R}^+)$ convergence

$$\phi_{\uparrow, t} \longrightarrow p_{\uparrow} \phi_{\infty}(a, v), \quad \phi_{\downarrow, t} \longrightarrow p_{\downarrow} \phi_{\infty}(a, v),$$

where $p_{\uparrow}$ and $p_{\downarrow}$ are probabilities that the averaged random walk exits above or below S respectively.

Moreover, the convergence is uniform for $|d| \leq \frac{1}{2}t^\gamma$.

Proof. Define the joint density $\Phi_{\uparrow, t}(a, v, w)$ for the position a , increment v for the over-jump of the boundary of S , and the size of the jump w which exits above S :

$$\Phi_{\uparrow, t}(a, v, w) = \sum_{n=1}^{\infty} \Pr[X_{t_n} = a, K_{t_n} = w, K_{t_n} - k - t^\gamma = v \geq 0, K_{t_m} \in S \text{ for } 0 \leq m < n] 1_{K_{t_n} > S}.$$

The definition for $\Phi_{\downarrow, t}(a, v, w)$ is analogous. Let primed densities be normalized (e.g. $\Phi'_{\uparrow, t} = p_{\uparrow}^{-1} \Phi_{\uparrow, t}$ and $\pi'_{\uparrow, t} = p_{\uparrow, t}^{-1} \pi_{\uparrow, t}$, where $p_{\uparrow, t}$ is the probability of leaving S from the top). We will show that $\Phi'_{\uparrow, t}$ converges in L^1 to

$$\Phi_{\infty}(a, v, w) = \pi_{\infty}(v, w) \frac{\frac{\kappa(a)}{\tilde{\kappa}} \mathcal{P}_a(w)}{\tilde{P}(w)}. \quad (\text{B.12})$$

Since $\phi'_{\uparrow, t}(a, v) = \int_{\mathbb{R}^+} dw \Phi_{\uparrow, t}(a, v, w)$, and $p_{\uparrow, t}$ converges to the probability that the random walk exits in the up direction by Proposition B.6, this would prove the result. In particular, since $|d| \leq \frac{1}{2}t^\gamma$ neither of the probabilities $p_{\infty}(\uparrow, t)$ or $p_{\infty}(\downarrow, t)$ will be close to zero. Also by the proof of B.6, the probability that the final momentum increment $w = K_{t_N} - K_{t_{N-1}}$ is greater than $2t^\gamma$, where t_N is the time of the last momentum jump leaving S and t_{N-1} is the time of previous momentum jump, decays as $O(t^{-2\gamma})$. This is true also for the random walk case. $\|\Phi_{\infty} \chi(w > 2t^\gamma)\|_1$ and $\|\Phi_{\uparrow, t} \chi(w > 2t^\gamma)\|_1$ thus vanish for large t .

Define $\|g\|_1^{(t)} = \|g \chi(w \leq 2t^\gamma, v < w)\|_1$. We placed the constraint $v < w$ in the indicator so that by Part (3) of Lemma B.1, $\pi'_{\infty}(v, w)$ is strictly positive and, in particular, we can divide by it. $\pi'_{\uparrow, t}(v, w)$ is also strictly positive for $v < w < 2t^\gamma$, since there is a non-zero density for jumping from k to $k + t^\gamma - w + v$ in one jump and then to $k + t^\gamma + v$ on a second jump.

We now focus on showing that $\|\Phi_{\infty} - \Phi_{\uparrow, t}\|_1^{(t)}$ tends to zero.

$$\frac{\Phi'_{\uparrow, t}(a, v, w)}{\pi'_{\uparrow, t}(v, w)} = \mathbb{E} \left[\Pr[X_{t_N} = a \mid K_{t_N} = t^\beta + t^\gamma + v, K_{t_N} - K_{t_{N-1}} = w, X_{t_{N-1}}] \right], \quad (\text{B.13})$$

since $\pi'_{\uparrow,t}(v, w) = \int_0^1 da \Phi'_{\uparrow,t}(a, v, w)$. We may write the conditional probability density in the expectation above as

$$\Pr [X_{t_N} = a \mid K_{t_N} = t^\beta + t^\gamma + v, K_{t_N} - K_{t_{N-1}} = w, X_{t_{N-1}}] = \frac{\mathcal{P}_a(w - \Delta(a)) \tilde{r}_{(X_{t_{N-1}}, K_{t_{N-1}})}(a)}{P_{(X_{t_{N-1}}, K_{t_{N-1}})}(w)}. \quad (\text{B.14})$$

$\Delta(a)$ is a drift term which was defined in Lemma B.5. $P_{(x,k)}(w)$ are defined as in the proof of Lemma B.5 as the difference in momentum (sum of one momentum jump plus some drift) between a starting time with state (x, k) and next time jump time. $\tilde{r}_{(x,k)}(a)$ is defined as in Lemma 5.2 as the distribution for the position of the particle at the next momentum jump.

Analogously to $\|g\|_1^{(t)}$ define the semi-norm $\|g\|_\infty^{(t)} = \|g\chi(|w| \leq 2t^\gamma, v < w)\|_\infty$.

Putting together (B.15)-(B.17),

$$\begin{aligned} \left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} \frac{\pi_\infty}{\Phi_\infty} - 1 \right\|_\infty^{(t)} &\leq \mathbb{E} \left[\sup_{w \geq 2t^\gamma, a} \left| \tilde{r}_{(X_{t_{N-1}}, K_{t_{N-1}})}(a) \frac{\bar{\kappa}}{\kappa(a)} \frac{\tilde{P}(w)}{P_{(X_{t_{N-1}}, K_{t_{N-1}})}(w)} - 1 \right| \right] \\ &\leq \sup_{y \geq t^\beta, a, x} \left[\tilde{r}_{(x,y)}(a) \frac{\bar{\kappa}}{\kappa(a)} \right] \mathbb{E} \left[\left| \frac{\tilde{P}(w)}{P_{(X_{t_{N-1}}, K_{t_{N-1}})}(w)} - 1 \right| \right] + \mathbb{E} \left[\sup_a \left| \tilde{r}_{(X_{t_{N-1}}, K_{t_{N-1}})}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \right] \\ &< (1 + 4\mathcal{R}\nu^{-2}t^{-\beta} + O(t^{-2\beta})) (2ct^{\gamma-\beta}) + 4\mathcal{R}\nu^{-2}t^{-\beta} + O(t^{-2\beta}) = O(t^{\gamma-2\beta}), \end{aligned}$$

where in the strict inequality we have used Part (2) Lemma 5.2 and Lemma B.5:

$$\sup_{0 \leq a \leq 1} \left| \tilde{r}_{(x,k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \leq 8\mathcal{R}\nu^{-2}t^{-\beta}, \quad \sup_{w \in R^+} \left| \frac{P_{(x,k)}(w)}{\tilde{P}(w)} - 1 \right| \leq 2ct^{\gamma-\beta}, \quad \text{for } |k| \geq \frac{1}{2}t^\beta$$

(where we doubled the constant factor in front of the higher order term to get rid of the lower), and we used that $|\frac{1}{b} - 1| \leq 2b$ for b in a small neighborhood around one.

By adding and subtracting by $\Phi'_{\uparrow,t} \frac{\pi'_{\uparrow,\infty}}{\pi'_{\uparrow,t}}$, and the triangle inequality

$$\begin{aligned} \|\Phi'_{\uparrow,t} - \Phi'_{\uparrow,\infty}\|_1^{(t)} &\leq \left\| \left(\frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} - \frac{\Phi'_{\uparrow,\infty}}{\pi'_{\uparrow,\infty}} \right) \pi'_{\uparrow,\infty} \right\|_1^{(t)} + \left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} (\pi'_{\uparrow,t} - \pi'_{\uparrow,\infty}) \right\|_1^{(t)} \\ &\leq \|\pi'_{\uparrow,\infty}\|_1 \left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} - \frac{\Phi'_{\uparrow,\infty}}{\pi'_{\uparrow,\infty}} \right\|_\infty^{(t)} + \|\pi'_{\uparrow,t} - \pi'_{\uparrow,\infty}\|_1 \left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} \right\|_\infty^{(t)}. \end{aligned}$$

$\|\pi'_{\uparrow,\infty}\|_1 = 1$ and so with the analysis above the left term on the second line tends to zero. For the right term, $\|\pi'_{\uparrow,t} - \pi'_{\uparrow,\infty}\|_1$ vanishes by Proposition B.6 and $\left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} \right\|_\infty^{(t)}$ is bounded by $\left\| \frac{\Phi_\infty}{\pi_\infty} \right\|_\infty^{(t)}$ plus a small number by the analysis above. $\square$

Proposition B.8 states the same results as in Proposition B.7 when there is some conditioning on the future of the particle's trajectories. For a particle starting with momentum in $(t^\beta, 2t^\beta)$, let $\theta_\uparrow$, $\theta_\downarrow$, and τ be the first times the particle has a momentum landing above $2t^\beta$, below $-2t^\beta$ and below t^β respectively. By the same argument as in Lemma 6.1, $\theta_\uparrow \wedge \theta_\downarrow$ has finite expectation.

Proposition B.8. Assume (i)-(ii) from List 2.2. Consider the dynamics conditioned to have momentum jumps bounded by $t^{\frac{\gamma}{2}}$. Let (X_s, K_s) evolve from an initial point (x, k) with $\frac{3}{2}t^\beta - t^{\frac{\gamma}{2}} \leq k \leq \frac{3}{2}t^\beta$. Let $S = \{y \in \mathbb{R} \mid k - t^\gamma < y < k + t^\gamma\}$ for $0 < \gamma < \frac{1}{4}\beta$.

The boundary crossing densities $\psi_{\uparrow, t}$, $\psi_{\downarrow, t}$ for trajectories that are conditioned so that $\tau < \theta_{\uparrow} < \theta_{\downarrow}$, have $L^1([0, 1] \times \mathbb{R}^+)$ convergence

$$\psi_{\uparrow, t} \longrightarrow \frac{1}{2} \phi_{\infty}, \quad \psi_{\downarrow, t} \longrightarrow \frac{1}{2} \phi_{\infty}.$$

The convergence is uniform for the allowed range of k . The same statements hold for $\psi_{\uparrow, t}$, $\psi_{\downarrow, t}$ for trajectories which are conditioned so that $\theta_{\downarrow} < \theta_{\uparrow}$.

Proof. Define the joint density $\Phi_{\uparrow, t}(a, v, w)$ for the position a , increment v for the over-jump of the boundary of S , and the size of the jump w which exits above S :

$$\Phi_{\uparrow, t}(a, v, w) = \sum_{n=1}^{\infty} \Pr[X_{t_n} = a, K_{t_n} = w, K_{t_n} - k - t^\gamma = v \geq 0, K_{t_m} \in S \text{ for } 0 \leq m < n] 1_{K_{t_n} > S}.$$

The definition for $\Phi_{\downarrow, t}(a, v, w)$ is analogous. Let primed densities be normalized (e.g. $\Phi'_{\uparrow, t} = p_{\uparrow}^{-1} \Phi_{\uparrow, t}$ and $\pi'_{\uparrow, t} = p_{\uparrow, t}^{-1} \pi_{\uparrow, t}$, where $p_{\uparrow, t}$ is the probability of leaving S from the top). We will show that $\Phi'_{\uparrow, t}$ converges in L^1 to

$$\Phi_{\infty}(a, v, w) = \pi_{\infty}(v, w) \frac{\frac{\kappa(a)}{\bar{\kappa}} \mathcal{P}_a(w)}{\tilde{P}(w)}. \quad (\text{B.15})$$

Since $\phi'_{\uparrow, t}(a, v) = \int_{\mathbb{R}^+} dw \Phi_{\uparrow, t}(a, v, w)$, and $p_{\uparrow, t}$ converges to the probability that the random walk exits in the up direction by Proposition B.6, this would prove the result. In particular, since $|d| \leq \frac{1}{2}t^\gamma$ neither of the probabilities $p_{\infty}(\uparrow, t)$ or $p_{\infty}(\downarrow, t)$ will be close to zero. Also by the proof of B.6, the probability that the final momentum increment $w = K_{t_N} - K_{t_{N-1}}$ is greater than $2t^\gamma$, where t_N is the time of the last momentum jump leaving S and t_{N-1} is the time of previous momentum jump, decays as $O(t^{-2\gamma})$. This is true also for the random walk case. $\|\Phi_{\infty} \chi(w > 2t^\gamma)\|_1$ and $\|\Phi_{\uparrow, t} \chi(w > 2t^\gamma)\|_1$ thus vanish for large t .

Define $\|g\|_1^{(t)} = \|g \chi(w \leq 2t^\gamma, v < w)\|_1$. We placed the constraint $v < w$ in the indicator so that by Part (3) of Lemma B.1, $\pi'_{\infty}(v, w)$ is strictly positive and, in particular, we can divide by it. $\pi'_{\uparrow, t}(v, w)$ is also strictly positive for $v < w < 2t^\gamma$, since there is a non-zero density for jumping from k to $k + t^\gamma - w + v$ in one jump and then to $k + t^\gamma + v$ on a second jump.

We now focus on showing that $\|\Phi_{\infty} - \Phi_{\uparrow, t}\|_1^{(t)}$ tends to zero.

$$\frac{\Phi'_{\uparrow, t}(a, v, w)}{\pi'_{\uparrow, t}(v, w)} = \mathbb{E} \left[\Pr[X_{t_N} = a \mid K_{t_N} = t^\beta + t^\gamma + v, K_{t_N} - K_{t_{N-1}} = w, X_{t_{N-1}}] \right], \quad (\text{B.16})$$

since $\pi'_{\uparrow, t}(v, w) = \int_0^1 da \Phi'_{\uparrow, t}(a, v, w)$. We may write the conditional probability density in the expectation above as

$$\begin{aligned} \Pr[X_{t_N} = a \mid K_{t_N} = t^\beta + t^\gamma + v, K_{t_N} - K_{t_{N-1}} = w, X_{t_{N-1}}] \\ = \frac{\mathcal{P}_a(w - \Delta(a)) \tilde{r}_{(X_{t_{N-1}}, K_{t_{N-1}})}(a)}{P_{(X_{t_{N-1}}, K_{t_{N-1}})}(w)}. \end{aligned} \quad (\text{B.17})$$

$\Delta(a)$ is a drift term which was defined in Lemma B.5. $P_{(x,k)}(w)$ are defined as in the proof of Lemma B.5 as the difference in momentum (sum of one momentum jump plus some drift) between a starting time with state (x, k) and next time jump time. $\tilde{r}_{(x,k)}(a)$ is defined as in Lemma 5.2 as the distribution for the position of the particle at the next momentum jump.

Analogously to $\|g\|_1^{(t)}$ define the semi-norm $\|g\|_\infty^{(t)} = \|g \chi(|w| \leq 2t^\gamma, v < w)\|_\infty$.

Putting together (B.15)-(B.17),

$$\begin{aligned} \left\| \frac{\Phi'_{\uparrow,t} \pi_\infty}{\pi'_{\uparrow,t} \Phi_\infty} - 1 \right\|_\infty^{(t)} &\leq \mathbb{E} \left[\sup_{w \geq 2t^\gamma, a} \left| \tilde{r}_{(X_{t_{N-1}}, K_{t_{N-1}})}(a) \frac{\bar{\kappa}}{\kappa(a)} \frac{\tilde{P}(w)}{P_{(X_{t_{N-1}}, K_{t_{N-1}})}(w)} - 1 \right| \right] \\ &\leq \sup_{y \geq t^\beta, a, x} \left[\tilde{r}_{(x,y)}(a) \frac{\bar{\kappa}}{\kappa(a)} \right] \mathbb{E} \left[\left| \frac{\tilde{P}(w)}{P_{(X_{t_{N-1}}, K_{t_{N-1}})}(w)} - 1 \right| \right] + \mathbb{E} \left[\sup_a \left| \tilde{r}_{(X_{t_{N-1}}, K_{t_{N-1}})}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \right] \\ &< (1 + 4\mathcal{R}\nu^{-2}t^{-\beta} + O(t^{-2\beta})) (2ct^{\gamma-\beta}) + 4\mathcal{R}\nu^{-2}t^{-\beta} + O(t^{-2\beta}) = O(t^{\gamma-2\beta}), \end{aligned}$$

where in the strict inequality we have used Part (2) Lemma 5.2 and Lemma B.5:

$$\sup_{0 \leq a \leq 1} \left| \tilde{r}_{(x,k)}(a) \frac{\bar{\kappa}}{\kappa(a)} - 1 \right| \leq 8\mathcal{R}\nu^{-2}t^{-\beta}, \quad \sup_{w \in R^+} \left| \frac{P_{(x,k)}(w)}{\tilde{P}(w)} - 1 \right| \leq 2ct^{\gamma-\beta}, \quad \text{for } |k| \geq \frac{1}{2}t^\beta$$

(where we doubled the constant factor in front of the higher order term to get rid of the lower), and we used that $|\frac{1}{b} - 1| \leq 2b$ for b in a small neighborhood around one.

By adding and subtracting by $\Phi'_{\uparrow,t} \frac{\pi'_{\uparrow,\infty}}{\pi'_{\uparrow,t}}$, and the triangle inequality

$$\begin{aligned} \|\Phi'_{\uparrow,t} - \Phi'_{\uparrow,\infty}\|_1^{(t)} &\leq \left\| \left(\frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} - \frac{\Phi'_{\uparrow,\infty}}{\pi'_{\uparrow,\infty}} \right) \pi'_{\uparrow,\infty} \right\|_1^{(t)} + \left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} (\pi'_{\uparrow,t} - \pi'_{\uparrow,\infty}) \right\|_1^{(t)} \\ &\leq \|\pi'_{\uparrow,\infty}\|_1 \left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} - \frac{\Phi'_{\uparrow,\infty}}{\pi'_{\uparrow,\infty}} \right\|_\infty^{(t)} + \|\pi'_{\uparrow,t} - \pi'_{\uparrow,\infty}\|_1 \left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} \right\|_\infty^{(t)}. \end{aligned}$$

$\|\pi'_{\uparrow,\infty}\|_1 = 1$ and so with the analysis above the left term on the second line tends to zero. For the right term, $\|\pi'_{\uparrow,t} - \pi'_{\uparrow,\infty}\|_1$ vanishes by Proposition B.6 and $\left\| \frac{\Phi'_{\uparrow,t}}{\pi'_{\uparrow,t}} \right\|_\infty^{(t)}$ is bounded by $\left\| \frac{\Phi_\infty}{\pi_\infty} \right\|_\infty^{(t)}$ plus a small number by the analysis above. $\square$

In the next corollary, like Proposition B.2, there is only one boundary. However, we place an optional time constraint ϱt , $\varrho \in (0, \infty]$ here on the amount of time the particle is allowed to have before reaching the boundary.

Corollary B.9. *Assume (i)-(ii) from List 2.2. Let (X_s, K_s) evolve according to the dynamics from some initial point (x, k) for $2t^\beta \leq k \leq 4t^\beta$, $\frac{1}{4} \leq \beta < \frac{1}{2}$. Define the density*

$$\phi_t(a, v) = \sum_{n=1}^{\infty} \Pr[X_{t_n} = a, t^\beta - K_{t_n} = v \geq 0, K_{t_m} > t^\beta \text{ for } 0 \leq m < n, t_n < \varrho t].$$

In the limit $t \rightarrow \infty$,

$$\phi_t(a, v) \rightarrow \phi_\infty(a, v).$$

The convergence is uniform over all starting points (x, k) .

Proof. First, we will argue that the probability $p_t = \|\phi_t\|_1$ that the particle jumps below t^β in the interval $[0, \varrho t]$ approaches one as $t \rightarrow \infty$ for $\varrho \in \mathbb{R}^+$.

$$\begin{aligned} p_t &= \Pr \left[\inf_{0 \leq r \leq \varrho t} t^{-\frac{1}{2}} K_r \geq t^{\beta-\frac{1}{2}} \right] \\ &\geq \Pr \left[t^{\beta-\frac{1}{2}} + t^{-\frac{1}{2}} k + \sup_{0 \leq r \leq \varrho t} \left| t^{-\frac{1}{2}} \int_0^r ds \frac{dV}{dx}(X_s) \right| \leq \sup_{0 \leq r \leq \varrho t} -t^{-\frac{1}{2}} M_r \right], \quad (\text{B.18}) \end{aligned}$$

since $K_r = k + M_r + \int_0^r ds \frac{dV}{dx}(X_s)$. By Chebyshev's inequality and that $k \leq 4t^\beta$.

$$\begin{aligned} \Pr \left[t^{\beta-\frac{1}{2}} + t^{-\frac{1}{2}} k + \sup_{0 \leq r \leq \varrho t} \left| t^{-\frac{1}{2}} \int_0^r ds \frac{dV}{dx}(X_s) \right| \geq \epsilon \right] \\ \leq \epsilon^{-2} \mathbb{E} \left[5t^{\beta-\frac{1}{2}} + \sup_{0 \leq r \leq \varrho t} \left| t^{-\frac{1}{2}} \int_0^r ds \frac{dV}{dx}(X_s) \right|^2 \right]. \quad (\text{B.19}) \end{aligned}$$

The right-side converges to zero by Lemma 5.3, and thus the probability that the supremum of the drift is greater than any finite ϵ_1 tends to zero.

By (3.1), $t^{-\frac{1}{2}} M_{st}$, $s \in [0, 1]$ converges to a Brownian motion $\mathbf{B}_s$. Since the supremum over an interval $[0, 1]$ is a uniformly continuous functional on elements $L^\infty[0, 1]$, $\sup_{0 \leq s \leq st} -t^{-\frac{1}{2}} M_{st}$ converges in distribution to $\sup_{0 \leq s \leq 1} -\mathbf{B}_s$. In the large t limit,

$$\Pr \left[\sup_{0 \leq s \leq \varrho t} -t^{-\frac{1}{2}} M_{st} > 2\epsilon \right] \longrightarrow \Pr \left[\sup_{0 \leq s \leq \varrho} -\mathbf{B}_s > 2\epsilon \right].$$

Thus we can pick ϵ so that $\Pr[\sup_{0 \leq s \leq 1} -\mathbf{B}_s > 2\epsilon]$ is close to one, and then pick t so that both $\Pr[\sup_{0 \leq s \leq \varrho t} -t^{-\frac{1}{2}} M_{st} > 2\epsilon]$ is close to $\Pr[\sup_{0 \leq s \leq \varrho} -\mathbf{B}_s > 2\epsilon]$ and (B.19) is close to zero. By the inclusion exclusion principle

$$p_t \geq 1 - 2 \Pr \left[\sup_{0 \leq s \leq \varrho t} -t^{-\frac{1}{2}} M_{st} > 2\epsilon \right] \wedge \Pr \left[t^{\beta-\frac{1}{2}} + t^{-\frac{1}{2}} k + \sup_{0 \leq r \leq \varrho t} \left| t^{-\frac{1}{2}} \int_0^r ds \frac{dV}{dx}(X_s) \right| \geq \epsilon \right],$$

and so p_t can be made arbitrarily close to 1.

Now we continue with showing L^1 convergence of ϕ_t to ϕ_∞ . Since $(1 - \|\phi_t\|)$ is negligible, it is sufficient to show $\phi_t \rightarrow (1 - \|\phi_t\|)\phi_\infty$. Let $\gamma < \frac{1}{3}\beta$. We construct a sequence of hitting times

$$\begin{aligned} \sigma_j &= \min\{s \in [0, \infty) \mid s > \theta_{j-1}, K_s < t^\beta + t^\gamma\}, \\ \theta_j &= \min\{s \in [0, \infty) \mid s > \sigma_j, K_s > t^\beta + 2t^\gamma\}. \end{aligned}$$

where we set $\theta_0 = 0$. Let N' be the number of σ_j 's in the interval $[0, \varrho t]$ before the first time t_N that K_{t_N} jumps below t^β . The process will usually have its first jump below t^β from a point with momentum in the interval $[t^\beta, t^\beta + \frac{1}{2}t^\gamma]$. In other words, $\Pr[|K_{t_{N-1}}| > t^\gamma]$ is small. If, as defined in Lemma B.4, $\pi_\downarrow(v)$ is the distribution for the over-jump for the lower boundary of the set $S = [t^\beta + t^\gamma, \infty)$ starting from some point in S , then by Lemma B.4 $\int_{\mathbb{R}^+} dv \pi_\downarrow(v) v^2 < J$. By Chebyshev's inequality

$$\Pr[|K_{t_{N-1}}| > t^\beta + \frac{1}{2}t^\gamma] \leq \int_{\mathbb{R}^+} dv \pi_\downarrow(v) \chi(|v| \geq \frac{1}{2}t^\gamma) \leq \frac{J}{4} t^{-2\gamma}.$$

The above comments allow us to write the following

$$\left\| \phi_t - \mathbb{E} \left[\sum_{j=1}^{\infty} \chi(N = j; K_{\sigma_j} - t^\beta \geq \frac{1}{2} t^\gamma) \frac{\phi_{\downarrow, t}^{(X_{\sigma_j}, K_{\sigma_j})}}{p(X_{\sigma_j}, K_{\sigma_j})} \right] \right\|_1 \leq \Pr [|K_{t_{N-1}}| > t^\gamma] \leq J t^\gamma,$$

where $\phi_{\downarrow, t}^{(X_{\sigma_j}, K_{\sigma_j})}$ is the lower boundary crossing distribution as in Proposition B.7 starting from the point $(X_{\sigma_j}, K_{\sigma_j})$, and $\int da dv \phi_{\downarrow, t}^{(X_{\sigma_j}, K_{\sigma_j})}(a, v) = p(X_{\sigma_j}, K_{\sigma_j})$ is the probability of exiting the domain $[t^\beta, t^\beta + 2t^\gamma)$ at the lower boundary. The difference d between the distance to the upper and lower boundaries $2t^\beta$ and $t^\beta + t^{2\gamma}$ respectively is $d = 2(t^\beta + t^\gamma - K_{\sigma_j})$.

$$\begin{aligned} & \left\| \mathbb{E} \left[\sum_{j=1}^{\infty} \chi(N = j; K_{\sigma_j} - t^\beta \geq \frac{1}{2} t^\gamma) \frac{\phi_{\downarrow, t}^{(X_{\sigma_j}, K_{\sigma_j})}}{p(X_{\sigma_j}, K_{\sigma_j})} \right] - p_t \pi_\infty \right\|_1 \\ & \leq \sum_{n=1}^{\infty} \Pr [\chi(N' = j)] \sup_{y - t^\beta \geq \frac{1}{2} t^\gamma, a} \left\| \frac{\phi_{\downarrow, t}^{(a, y)}}{p(a, y)} - \pi_\infty \right\|_1 \leq \sup_{y - t^\beta \geq \frac{1}{2} t^\gamma, a} \left\| \frac{\phi_{\downarrow, t}^{(a, y)}}{p(a, y)} - \pi_\infty \right\|_1 \end{aligned}$$

By Proposition B.7 $p(a, y)$, $y \geq t^\beta + \frac{1}{2} t^\gamma$ converge uniformly to the the probabilities $p_\downarrow$ for the averaged random walk to exit in down direction. $p_\downarrow$ are bounded away from zero, since the ratio of the distance to the lower boundary to the upper boundary is less than or equal to 3. Moreover, by Proposition B.9 $\phi_{\downarrow, t}^{(a, y)}$ converges uniformly to $p_\downarrow \pi_\infty$. □

References

- [1] G. Basile and A. Bovier: *Convergence of a kinetic equation to a fractional diffusion equation*, arXiv:0909.3385
- [2] F. Bonetto, J.L. Lebowitz, and L. Rey-Bellet: *Fourier's Law: a Challenge for Theorists*, Mathematical Physics 2000, Edited by A. Fokas, A. Grigoryan, T. Kibble and B. Zegarlinsky, Imperial College Press, 128-151 (2000).
- [3] K.L. Chung: *A Course in Probability Theory*, Academic Press, New York 1976.
- [4] A. De Masi, P.A. Ferrari, S. Goldstein, and D.W. Wick: *An Invariance Principle for Reversible Markov Processes. Applications to Random Motions in Random Environments*, J. Stat. Phys. **55**, 767–855 (1989).
- [5] S. Goldstein: *Antisymmetric functionals of reversible Markov processes*, Ann. Inst. Henri Poincaré **31**, 177–190 (1995).
- [6] M. Hairer, G.A. Pavliotis: *From ballistic to diffusive behavior in periodic potentials*, J. Stat. Phys. **131**, 175-202 (2008).
- [7] P. Hall, C.C. Heyde *Martingale Limit Theory and Its Application*, Academic Press, 1980.
- [8] D.V. Husak: *On the joint distribution of the time and value of the first overjump for homogeneous processes with independent increments* Teor. Ver. Primen. **14**, 15-23 (1969).

- [9] M. Jara and T. Komorowski: *Non-Markovian limits of additive functionals of Markov processes*, `arXiv:0905.2163`.
- [10] M. Jara, T. Komorowski, S. Olla: *Limit theorems for additive functionals of a Markov chain*, *Ann. Appl. Probab.* **19**, 2270-2300 (2009).
- [11] I. Karatzas, S.E. Shreve: *Brownian Motion and Stochastic Calculus*, Springer-Verlag, 1988.
- [12] C. Kipnis and S.R.S. Varadhan: *Central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusions*, *Commun. Math. Phys.* **104**, 1–19 (1986).
- [13] T. Komorowski, C. Landim and S. Olla: *Fluctuations in Markov Processes*. See webpage <http://w3.impa.br/~landim/notas.html>.
- [14] D. Pollard: *Convergence of Stochastic Processes*, Springer-Verlag, 1984.
- [15] D. Weihs, M.A. Teitell and T.G. Mason: *Simulations of complex particle transport in heterogeneous active liquids*, *Microfluid Nanofluid* **3**, 227–237 (2007).